

Aspects of entanglement entropy for gauge theories

Ronak M. Soni and Sandip P. Trivedi

*Department of Theoretical Physics, Tata Institute of Fundamental Research,
Dr. Homi Bhabha Road, Colaba, Mumbai, 400005, India*

E-mail: ronak@theory.tifr.res.in, sandip@theory.tifr.res.in

ABSTRACT: A definition for the entanglement entropy in a gauge theory was given recently in arXiv:1501.02593. Working on a spatial lattice, it involves embedding the physical state in an extended Hilbert space obtained by taking the tensor product of the Hilbert space of states on each link of the lattice. This extended Hilbert space admits a tensor product decomposition by definition and allows a density matrix and entanglement entropy for the set of links of interest to be defined. Here, we continue the study of this extended Hilbert space definition with particular emphasis on the case of Non-Abelian gauge theories.

We extend the electric centre definition of Casini, Huerta and Rosabal to the Non-Abelian case and find that it differs in an important term. We also find that the entanglement entropy does not agree with the maximum number of Bell pairs that can be extracted by the processes of entanglement distillation or dilution, and give protocols which achieve the maximum bound. Finally, we compute the topological entanglement entropy which follows from the extended Hilbert space definition and show that it correctly reproduces the total quantum dimension in a class of Toric code models based on Non-Abelian discrete groups.

KEYWORDS: Gauge Symmetry, Lattice Quantum Field Theory

ARXIV EPRINT: [1510.07455](https://arxiv.org/abs/1510.07455)

Contents

1	Introduction	1
2	The definition	4
2.1	General discussion	5
2.2	Non-Abelian theories	7
3	The electric centre definition	9
3.1	General discussion	10
3.2	Non-Abelian case	11
3.3	Entanglement entropy	13
4	Entanglement distillation and dilution	14
4.1	A toy model	15
4.2	The distillation protocol	18
4.3	The dilution protocol	22
4.4	Concluding comments on the toy model	25
4.5	Comments on gauge theories	26
5	Degeneracy of the non-Abelian toric code	28
5.1	The different superselection sectors	31
6	Entanglement entropy in the strong-coupling expansion	34
7	Conclusions	39

1 Introduction

Quantum systems are known to behave in essentially different ways from their classical counterparts. Entanglement is a key property that characterises quantum correlations and entanglement entropy provides a quantitative measure of some of these essential differences. For a bipartite system consisting of two parts A, B , with Hilbert spaces, $\mathcal{H}_A, \mathcal{H}_B$, respectively, which is in a pure state, $|\psi\rangle$, the entanglement entropy S_{EE} is obtained by constructing the density matrix of one of the two parts, say A , after tracing over the second one, B ,

$$\rho_A = \text{Tr}_{\mathcal{H}_B} |\psi\rangle\langle\psi| \quad (1.1)$$

and computing its von Neumann entropy

$$S_{EE} = -\text{Tr}_{\mathcal{H}_A} \rho_A \log \rho_A. \quad (1.2)$$

The Hilbert space of the full system in this case is given by the tensor product

$$\mathcal{H} = \mathcal{H}_A \otimes \mathcal{H}_B. \quad (1.3)$$

In a gauge theory the definition of the entanglement gets more complicated. It turns out that the Hilbert space of physical states, i.e., gauge-invariant states, does not admit a tensor product decomposition of the type eq. (1.3) in terms of the Hilbert space of states in a region A and its complement. Thus, it is not clear how to compute the entanglement described above. This is a very general feature in gauge theories and it is related to the presence of non-local operators like Wilson lines, which create non-local excitations, in these theories.

Several approaches have been suggested in the literature, see, [1–11], to circumvent this problem and arrive at a satisfactory definition of entanglement entropy for a gauge theory. Here we will follow the approach discussed in [1] (GST), see also [9], which is also related to the earlier work, [3, 4]. In this approach, by working on a spatial lattice, a definition of entanglement entropy for a gauge theory is given as follows. One first embeds the space of gauge-invariant states in a bigger space \mathcal{H} obtained by taking the tensor product of the Hilbert spaces on each link of the lattice. This space, \mathcal{H} , by definition admits a tensor product decomposition in terms of the Hilbert spaces for the set of links of interest, which we denote as \mathcal{H}_{in} , and the Hilbert space for the rest of the links, \mathcal{H}_{out} . We can write, $\mathcal{H} = \mathcal{H}_{\text{in}} \otimes \mathcal{H}_{\text{out}}$, which is analogous to eq. (1.3). It is then straightforward to obtain the density matrix, ρ_{in} , eq. (1.1), by taking a trace over \mathcal{H}_{out} , and from it the entropy, eq. (2.4). The definition given in GST works for any set of links, including in particular a set of links which are adjacent or close to each other in space. We will refer to it as the extended Hilbert space definition below.

This definition has several nice features. It is gauge-invariant. It meets the condition of strong subadditivity. And it can be shown to agree with a path integral definition of entanglement which follows from the replica trick.¹ The definition applies to both Abelian and Non-Abelian theories and can be readily extended to gauge theories with matter.

The purpose of this paper is to explore the extended Hilbert space definition further and elucidate some of its properties.

A different approach for defining the entanglement entropy was given in the work of Casini, Huerta and Rosabal (CHR) [2], see also [6]. In this approach one works with gauge-invariant states, and the algebra of operators acting in the region A and its complement B . For a gauge theory it was argued that this algebra has a non-trivial centre, and this centre is responsible for the space of gauge-invariant states not having a tensor product decomposition, eq. (1.3). One can overcome this obstacle by diagonalising the centre and going to sectors where it takes a fixed value. The Hilbert space of gauge-invariant states in each sector then does admit a tensor product decomposition, leading to a definition of the density matrix and the entanglement entropy. The different sectors are in fact superselection sectors since no local gauge-invariant operators, acting on either the inside

¹More accurately, since there can be ambiguities in the replica trick on the lattice, the extended Hilbert space definition agrees with the path integral implementing a particular version of the replica trick.

or the outside, can change the sector. It turns out that different choices of the centre are possible and give different definitions for the entanglement on the lattice. A particular choice, called the electric centre, corresponds to specifying the total electric flux entering the links of interest at each vertex lying on the boundary. By Gauss' law this is also equal, up to a sign, to the total flux leaving the links of interest at the boundary vertices. This choice leads to the electric centre definition of the entanglement entropy.

For Abelian theories it was shown that the extended Hilbert space definition agreed with the electric centre definition, [1, 2]. The extended Hilbert space definition can also be expressed as a sum over sectors carrying different electric fluxes into the region of interest, and the contributions from each such sector agrees with what arises in the electric centre definition for the Abelian case. As mentioned above, the extended Hilbert space definition also works for Non-Abelian theories. In this paper we show how the resulting entanglement entropy in the Non-Abelian case can also be expressed as a sum over different sectors with each sector specifying, in a gauge-invariant way, the electric flux entering the inside links. We then try to extend the electric centre definition, by analogy from the Abelian case, to the Non-Abelian case, and again obtain a result which is a sum over the different electric flux sectors. However, interestingly, we find that the two definitions lead to different contributions in each sector, resulting in a different final result for the entanglement entropy.

The difference between the two definitions is tied to an interesting subtlety which arises in the Non-Abelian case. In both the Abelian and Non-Abelian cases the entanglement entropy gets contributions of two kinds. One kind is a sort of “classical” term which arises due to the system having a probability for being in the different superselection sectors. The second kind is due to the entanglement entropy within each superselection sector. In the Non-Abelian case, in the extended Hilbert space definition, the classical term in turn has two contributions. These give rise to the first two terms in eq. (2.29), and eq. (2.30), which will be discussed in section 3 in more length. One contribution is analogous to the Abelian case and arises due to the probability of being in the different superselection sectors. The other contribution, which is intrinsically a feature of the Non-Abelian case, arises as follows. In sectors which carry non-trivial electric flux at a boundary vertex, the inside and outside links which meet at this vertex by themselves transform non-trivially under gauge transformations, but together must combine to be gauge-invariant. Since in the Non-Abelian case representations of the symmetry group are of dimension greater than one this results in non-trivial entanglement between the inside and outside links and the resulting additional contribution to the entropy. This contribution was also discussed in [4] and [8, 9]. It turns out that this contribution of the classical kind is absent in the electric centre definition.

In quantum information theory an operational measure of entanglement entropy is provided by comparing the entanglement of a bipartite system with a set of Bell pairs. This comparison is done by the process of entanglement distillation or dilution. It is well known that for a system with localised degrees of freedom, like a spin system or a scalar field theory, the resulting number of Bell pairs produced in distillation or consumed in dilution, agrees with the definition given above, eq. (2.4). More correctly, this is true in the asymptotic sense, when one works with n copies of the system and takes the $n \rightarrow \infty$ limit. Only local operations and classical communication (LOCC) are allowed in these processes.

It was already mentioned in [1], see also [2], that for gauge theories the extended Hilbert space definition is not expected to agree with such an operational definition. In particular the total entanglement extracted in the form of Bell pairs in distillation or dilution should be smaller. The difference arises because physical LOCC operations can only involve local gauge-invariant operations and these are insufficient to extract all the entropy. In this paper we also explore this question in greater depth. As was mentioned above, the entanglement entropy in the extended Hilbert space definition gets contributions of two kinds, a classical contribution, and a quantum contribution due to entanglement within each superselection sector. We show that in the asymptotic sense mentioned above, all the entanglement of the second kind can be extracted in distillation or dilution, but none of the contribution of the first kind is amenable to such extraction. We make this precise by presenting protocols which achieve the maximum number of extracted Bell pairs for distillation and dilution. This result is not unexpected, and was anticipated in the Abelian case, [2], since local gauge-invariant operators cannot change the superselection sectors and therefore are not expected to be able to access the entropy contribution of the first kind.

Since the full entanglement entropy in the extended Hilbert space definition cannot be operationally measured one might wonder whether the correct definition of entanglement is obtained by simply keeping that part which does agree with the measurements. It has been shown that the entanglement entropy can be useful for characterising systems with non-trivial quantum correlations in states with a mass gap. For example, the entanglement entropy has been used to define the topological entanglement entropy [12, 13] which in turn is related to important properties of the system like the degeneracy of ground states on a manifold of non-trivial topology, and the total quantum dimension of excitations in the system. In section 5 we calculate the topological entanglement entropy for a class of Toric code models [14] based on non-Abelian discrete groups using the extended Hilbert space definition. We show that the correct result is obtained. This result gets contributions from both the classical and non-classical terms in general. In the contribution from classical terms both the first and second terms in eq. (2.30) which were mentioned a few paragraphs above contribute. Thus dropping these terms, which cannot be measured in distillation or dilution, would not lead to the correct answers for various physical questions tied to entanglement in these systems.

This paper is organised as follows. We discuss the extended Hilbert space definition in section 2, then compare to the electric centre definition which is given for the Non-Abelian case in section 3. A discussion of the entanglement distillation and dilution protocols follows in section 4. An analysis of the Toric code models is presented in section 5. We end in section 6 with a discussion, for good measure, of the entanglement in the ground state of an $SU(2)$ gauge theory to first non-trivial order in the strong coupling expansion. We find that to this order the entanglement arises solely due to the classical terms.

2 The definition

In this section we review the extended Hilbert space definition of entanglement entropy given in GST, see also [9].

2.1 General discussion

We work in a lattice gauge theory in the Hamiltonian framework. The degrees of freedom of the theory live on links L_{ij} of a spatial lattice. Gauge transformations G_{V_i} are defined on vertices, V_i . Physical states are gauge-invariant and satisfy the condition

$$G_{V_i} |\psi\rangle = |\psi\rangle. \quad (2.1)$$

These physical states span the Hilbert space of gauge-invariant states, denoted by $\mathcal{H}_{\text{ginv}}$.

An extended Hilbert space is defined as follows. The degrees of freedom on each link form a Hilbert space \mathcal{H}_{ij} . The extended Hilbert space is then given by $\mathcal{H} = \otimes \mathcal{H}_{ij}$ where the tensor product is taken over all links. We are interested in defining the entanglement entropy of a set of links, which we loosely call the “inside links.” The remaining links, not in the inside, are the “outside links.” In figure 1, which shows a square lattice for example, the inside links are shown as solid lines, while the outside links are shown as dashed lines.

\mathcal{H} admits a tensor product decomposition in terms of the Hilbert space of the links of interest, \mathcal{H}_{in} and its complement, which is the Hilbert space of outside links, \mathcal{H}_{out} . Also, if $\mathcal{H}_{\text{ginv}}^\perp$ is the orthogonal complement of $\mathcal{H}_{\text{ginv}}$ then \mathcal{H} can be written as the sum

$$\mathcal{H} = \mathcal{H}_{\text{ginv}} \oplus \mathcal{H}_{\text{ginv}}^\perp. \quad (2.2)$$

To define the entropy we regard $|\psi\rangle \in \mathcal{H}$, then trace over \mathcal{H}_{out} to get the density matrix

$$\rho_{\text{in}} = \text{Tr}_{\mathcal{H}_{\text{out}}} |\psi\rangle\langle\psi|. \quad (2.3)$$

The entanglement entropy is then defined as

$$S_{EE} = -\text{Tr}_{\mathcal{H}_{\text{in}}} \rho_{\text{in}} \log \rho_{\text{in}}. \quad (2.4)$$

This definition has several nice properties that were reviewed in GST. It is unambiguous and gauge-invariant. And it meets the strong subadditivity condition. We also note that the density matrix ρ_{in} correctly gives rise to the expectation value of any gauge-invariant operator \hat{O} which acts on the inside links. This is because $|\psi\rangle$ is orthogonal to all states in $\mathcal{H}_{\text{ginv}}^\perp$.

S_{EE} can be expressed in a manifestly gauge-invariant manner in terms of superselection sectors as follows. We define inside vertices as those on which only links securely in the inside set end, outside vertices as those on which only outside links end, and boundary vertices as those on which some links inside and some outside links end. In figure 1, for example, V_1 is a boundary vertex on which three inside links and one outside link end. A superselection sector is specified by a vector \mathbf{k} which gives the total electric flux entering the inside links at all the boundary vertices. Labelling the boundary vertices as $V_i, i = 1, \dots, N$, the i^{th} entry k_i of \mathbf{k} is the total electric flux entering the inside at the V_i boundary vertex. Then the inside Hilbert space \mathcal{H}_{in} can be written as a sum

$$\mathcal{H}_{\text{in}} = \oplus \mathcal{H}_{\text{in}}^{\mathbf{k}}, \quad (2.5)$$

where $\mathcal{H}_{\text{in}}^{\mathbf{k}}$ is the subspace of \mathcal{H}_{in} corresponding to the sector with flux \mathbf{k} .




Figure 1. An example of a region in a lattice. The solid lines are the inside links, and the dotted lines are outside links. V_1 is an example of what we call a boundary vertex, one which has both inside and outside links ending on it.

It can be shown that ρ_{in} is block diagonal in the different sectors and takes the form

$$\rho_{\text{in}} = \oplus \rho_{\text{in}}^{\mathbf{k}}, \quad (2.6)$$

where $\rho_{\text{in}}^{\mathbf{k}}$ acts on $\mathcal{H}_{\text{in}}^{\mathbf{k}}$.

S_{EE} can then be expressed as

$$S_{EE} = - \sum_{\mathbf{k}} \text{Tr}_{\mathcal{H}_{\text{in}}^{\mathbf{k}}} \rho_{\text{in}}^{\mathbf{k}} \log \rho_{\text{in}}^{\mathbf{k}}. \quad (2.7)$$

Let

$$\rho_{\text{in}}^{\mathbf{k}} = p_{\mathbf{k}} \tilde{\rho}_{\text{in}}^{\mathbf{k}}, \quad (2.8)$$

where

$$\text{Tr}_{\mathcal{H}_{\text{in}}^{\mathbf{k}}} \rho_{\text{in}}^{\mathbf{k}} = p_{\mathbf{k}} \quad (2.9)$$

so that

$$\text{Tr}_{\mathcal{H}_{\text{in}}^{\mathbf{k}}} \tilde{\rho}_{\text{in}}^{\mathbf{k}} = 1. \quad (2.10)$$

Then eq. (2.7) gives

$$S_{EE} = - \sum_{\mathbf{k}} p_{\mathbf{k}} \log p_{\mathbf{k}} - \sum_{\mathbf{k}} p_{\mathbf{k}} \text{Tr}_{\mathcal{H}_{\text{in}}^{\mathbf{k}}} \tilde{\rho}_{\text{in}}^{\mathbf{k}} \log \tilde{\rho}_{\text{in}}^{\mathbf{k}}. \quad (2.11)$$

The different sectors specified by the electric flux \mathbf{k} are called superselection sectors because no gauge-invariant operator acting on the inside links alone can change these sectors. On the other hand gauge-invariant operators which act on both inside and outside links can change these sectors, e.g., Wilson loops that extend from the inside to the outside links.

The description above applies to all gauge theories without matter. These include discrete Abelian theories, like a \mathbb{Z}_2 theory, continuous Abelian theories, e.g., a $U(1)$ theory, and finally Non-Abelian gauge theories, like an $SU(2)$ gauge theory.

2.2 Non-Abelian theories

For the Abelian cases a specification of the flux k_i at each boundary vertex is straightforward. On the other hand, for the Non-Abelian case there is an important subtlety which we now discuss. This subtlety has to do with the fact that the irreducible representations in the Non-Abelian case are not one-dimensional.

Consider as a specific example the $SU(2)$ group, where each link, L_{ij} , has as its degree of freedom a 2×2 matrix $U_{ij} \in SU(2)$. The corresponding state in the Hilbert space \mathcal{H}_{ij} is denoted by $|U_{ij}\rangle$. There are two sets of operators $\hat{J}_{ij}^a, a = 1, 2, 3$, and $\hat{\mathcal{J}}_{ij}^a, a = 1, 2, 3$, which acts as generators of the algebra, acting on the left and right respectively,

$$e^{i\epsilon^a \hat{J}_{ij}^a} |U_{ij}\rangle = \left| \left(1 + i\epsilon^a \frac{\sigma^a}{2} \right) U_{ij} \right\rangle \quad (2.12)$$

$$e^{i\epsilon^a \hat{\mathcal{J}}_{ij}^a} |U_{ij}\rangle = \left| U_{ij} \left(1 - i\epsilon^a \frac{\sigma^a}{2} \right) \right\rangle. \quad (2.13)$$

The links L_{ij} are oriented to emanate from vertex V_i and end on V_j . It is easy to see that with the definitions above

$$\hat{\mathcal{J}}_{ij}^a = \hat{J}_{ji}^a, a = 1, 2, 3. \quad (2.14)$$

Also, one can show that,

$$(\hat{J}_{ij})^2 = (\hat{\mathcal{J}}_{ij})^2. \quad (2.15)$$

The gauge transformation at V_i is then generated by

$$G_{V_i} = e^{i\epsilon^a \sum_j \hat{J}_{ij}^a}, \quad (2.16)$$

where the sum on the r.h.s. is over all links emanating from V_i .

Now consider a boundary vertex, V_i . An example is the vertex V_1 shown in figure 1. The total electric flux carried to outside from V_i is given by

$$\hat{J}_{T,\text{out},i}^a = \sum_j \hat{J}_{ij}^a, \quad (2.17)$$

where the sum is over all links which are in the outside set. Similarly the total flux carried inside is given by

$$\hat{J}_{T,\text{in},i}^a = \sum_j \hat{J}_{ij}^a, \quad (2.18)$$

with the sum being over all links going in.

Gauss' law says that

$$\hat{J}_{T,\text{out},i}^a = -\hat{J}_{T,\text{in},i}^a. \quad (2.19)$$

The superselection sectors are then specified by a choice of $k_i = (\hat{J}_{T,\text{in},i})^2 = (\hat{J}_{T,\text{out},i})^2$ for all boundary vertices V_i . Since the operators, $(\hat{J}_{T,\text{in},i})^2, (\hat{J}_{T,\text{out},i})^2$ are gauge-invariant, this is a gauge-invariant characterisation of the superselection sectors.

Now we come to the subtlety. It is easy to see that the density matrix which arises for a gauge-invariant state by tracing over \mathcal{H}_{out} satisfies the relation

$$[\rho_{\text{in}}, \hat{J}_{T,\text{in},i}^a] = 0. \quad (2.20)$$

Similarly, all gauge-invariant operators acting on inside links must also commute with $\hat{J}_{T,\text{in},i}^a$.

Now suppose

$$k_i = j_i(j_i + 1), \quad (2.21)$$

corresponding to the $2j_i + 1$ dimensional irreducible representation of $\text{SU}(2)$. Then $\mathcal{H}_{\text{in}}^{\mathbf{k}}$ must furnish a corresponding representation of $\hat{J}_{T,\text{in},i}^a$. These two features mean that $\mathcal{H}_{\text{in}}^{\mathbf{k}}$ can be written as a tensor product

$$\mathcal{H}_{\text{in}}^{\mathbf{k}} = \mathcal{H}^1 \otimes \mathcal{H}^2 \cdots \otimes \mathcal{H}^i \otimes \cdots \otimes \mathcal{H}^n \otimes \hat{\mathcal{H}}_{\text{in}}^{\mathbf{k}}, \quad (2.22)$$

where \mathcal{H}^i is a $2j_i + 1$ dimensional Hilbert space and $i = 1 \dots n$ label the boundary vertices. All gauge-invariant operators, and also $\rho_{\text{in}}^{\mathbf{k}}$, act non-trivially only on the last term in the tensor product, $\hat{\mathcal{H}}_{\text{in}}^{\mathbf{k}}$, and are trivial on the other factors. E.g., $\rho_{\text{in}}^{\mathbf{k}}$ can be written as

$$\rho_{\text{in}}^{\mathbf{k}} = \mathbf{I}_{2j_1+1} \otimes \cdots \otimes \mathbf{I}_{2j_i+1} \otimes \cdots \otimes \mathbf{I}_{2j_n+1} \otimes \hat{\rho}_{\text{in}}^{\mathbf{k}}, \quad (2.23)$$

where $\hat{\rho}_{\text{in}}^{\mathbf{k}}$ acts on $\hat{\mathcal{H}}_{\text{in}}^{\mathbf{k}}$, and \mathbf{I}_{2j_i+1} denotes the identity operator acting on \mathcal{H}^i . A gauge-invariant operator \mathcal{O} acting on inside links alone cannot change the superselection sectors. In the sector \mathbf{k} this takes the form

$$\mathcal{O}^{\mathbf{k}} = \mathbf{I}_{2j_1+1} \otimes \cdots \otimes \mathbf{I}_{2j_i+1} \otimes \cdots \otimes \mathbf{I}_{2j_n+1} \otimes \hat{\mathcal{O}}^{\mathbf{k}}, \quad (2.24)$$

where $\hat{\mathcal{O}}^{\mathbf{k}}$ acts on $\hat{\mathcal{H}}_{\text{in}}^{\mathbf{k}}$.

On the other hand, $\hat{J}_{T,\text{in},i}^a$, $a = 1, 2, 3$, act non-trivially only on \mathcal{H}^i and not on the other factors,

$$\hat{J}_{T,\text{in},i}^a = \mathbf{I}_{2j_1+1} \otimes J_i^a \otimes \cdots \otimes \mathbf{I}_{2j_n+1} \otimes \mathbf{I}, \quad (2.25)$$

where J_i^a denotes a matrix in the $2j_i + 1$ dimensional representation of $\text{SU}(2)$ acting on the \mathcal{H}^i Hilbert space.

As a result the second term in eq. (2.11) can be further split in two. The normalised density matrix, $\tilde{\rho}_{\text{in}}^{\mathbf{k}}$ was defined in eq. (2.8), eq. (2.10), above by rescaling the density matrix. This operator can also be written in the form eq. (2.23),

$$\tilde{\rho}_{\text{in}}^{\mathbf{k}} = \mathbf{I}_{2j_1+1} \otimes \cdots \otimes \mathbf{I}_{2j_i+1} \otimes \cdots \otimes \mathbf{I}_{2j_n+1} \otimes \hat{\tilde{\rho}}_{\text{in}}^{\mathbf{k}} \quad (2.26)$$

with $\hat{\tilde{\rho}}_{\text{in}}^{\mathbf{k}}$ replacing $\hat{\rho}_{\text{in}}^{\mathbf{k}}$. It is now convenient to further rescale and define the operator

$$\bar{\rho}_{\text{in}}^{\mathbf{k}} = \left(\prod_i (2j_i + 1) \right) \hat{\tilde{\rho}}_{\text{in}}^{\mathbf{k}}. \quad (2.27)$$

It is easy to see from eq. (2.23) that this operator satisfies the condition

$$\text{Tr}_{\hat{\mathcal{H}}_{\text{in}}^{\mathbf{k}}} \bar{\rho}_{\text{in}}^{\mathbf{k}} = 1. \quad (2.28)$$

The entanglement entropy can then be written as a sum of three terms

$$S_{EE} = - \sum_{\mathbf{k}} p_{\mathbf{k}} \log p_{\mathbf{k}} + \sum_{\mathbf{k}} p_{\mathbf{k}} \left(\sum_i \log(2j_i + 1) \right) - \sum_{\mathbf{k}} p_{\mathbf{k}} \text{Tr}_{\hat{\mathcal{H}}_{\text{in}}^{\mathbf{k}}} \bar{\rho}_{\text{in}}^{\mathbf{k}} \log \bar{\rho}_{\text{in}}^{\mathbf{k}}. \quad (2.29)$$

Here \mathbf{k} denotes the different superselection sectors. The sum over index i in the second term is over all boundary vertices, with j_i being the value of the total incoming angular momentum, $(\hat{J}_{T,\text{in}}^a)^2$ at the i^{th} boundary vertex, related to k_i by eq. (2.21). Note that the last trace is over the $\hat{\mathcal{H}}_{\text{in}}^{\mathbf{k}}$ subspace.

The second term in eq. (2.29) then is the extra contribution which arises in the Non-Abelian case since the irreducible representations can have dimension greater than unity. Physically, the total angular momentum at a boundary vertex must sum to zero. This results in a correlation between the in-going and out-going links at the vertex. The in-going links together give rise to a state in the $2j_i + 1$ dimension representation and so do the out-going links. Then these two states combine to give a singlet of the total angular momentum emanating from the vertex. The requirement of being in a singlet state entangles the state of the ingoing and outgoing states non-trivially, since the representations are not one dimensional. This extra entanglement results in the second term in eq. (2.29).

We have chosen the $SU(2)$ example for simplicity. The generalisation to any Non-Abelian group is immediate. The analogue of $\hat{J}_{T,\text{in},i}^a, \hat{J}_{T,\text{out},i}^a$ are the generators of the group acting on the ingoing and outgoing links. Superselection sectors are specified by giving the irreducible representations, r_i , under which these generators transform, at all boundary vertices,² V_i . And in the second term in eq. (2.29), $\log(2j_i + 1)$ is replaced by $\log d_i$ where d_i is the dimension of the r_i representation. As a result the entanglement is given by

$$S_{EE} = - \sum_{\mathbf{k}} p_{\mathbf{k}} \log p_{\mathbf{k}} + \sum_{\mathbf{k}} p_{\mathbf{k}} \left(\sum_i \log d_i \right) - \sum_{\mathbf{k}} p_{\mathbf{k}} \text{Tr}_{\hat{\mathcal{H}}_{\text{in}}^{\mathbf{k}}} \bar{\rho}_{\text{in}}^{\mathbf{k}} \log \bar{\rho}_{\text{in}}^{\mathbf{k}}. \quad (2.30)$$

We end with some concluding comment. The first term in eq. (2.11) for the Abelian case is sometimes referred to as a kind of “classical” contribution to the entropy. It is the minimum entropy which would arise from a density matrix, eq. (2.6), which meets the condition, eq. (2.9). This minimum contribution would arise if $\bar{\rho}^{\mathbf{k}}$, eq. (2.8), in each superselection sector corresponds to a pure state. Similarly, we may regard the first two terms in eq. (2.30) for the non-Abelian case also as a classical contribution. It too is the minimum contribution which arises for a density matrix which satisfies the condition, eq. (2.9), and which has the form eq. (2.23) required by eq. (2.20).

Gauge-invariant operators acting on only the inside or outside links cannot change the superselection sectors labelled by \mathbf{k} , and the probabilities $p_{\mathbf{k}}$ or the values of d_i . This suggests that the minimal contribution to the entanglement represented by the first term in eq. (2.11) and the first two terms in eq. (2.30) cannot be extracted in dilution or distillation experiments. We will see in section 4 that this expectation is indeed borne out.

3 The electric centre definition

For the Abelian case a different approach for defining the entanglement entropy using only the gauge-invariant Hilbert space of states $\mathcal{H}_{\text{ginv}}$ and not the extended Hilbert space was adopted in [2], see also, [6].

²More correctly if the outgoing links transform as the r_i representation then the ingoing links must transform as the conjugate representation \bar{r}_i . Together these give the identity in a unique way.

It was shown in [2] and [1] that the electric centre choice of CHR leads to a definition of entanglement entropy that coincides with the extended Hilbert space definition given above in the Abelian case. Here we examine the electric centre definition in the Non-Abelian case and find that it in fact differs from the definition given above, precisely in the second term on the r.h.s. of eq. (2.30).

3.1 General discussion

The essential point in the definition given by CHR is to focus on the algebra of gauge-invariant operators, \mathcal{A}_{in} , and \mathcal{A}_{out} , which act on the inside and outside links respectively. These algebras, it was argued, have a non-trivial centre,

$$\mathcal{A}_{\text{in}} \cap \mathcal{A}_{\text{out}} = \mathcal{Z}, \quad (3.1)$$

for a gauge theory, which commutes with both \mathcal{A}_{in} and \mathcal{A}_{out} . This non-trivial centre is the essential obstacle to obtaining a tensor product decomposition of $\mathcal{H}_{\text{ginv}}$.

To circumvent this problem, in the definition given by CHR, one works in sectors where the centre is diagonalised. Within each sector, labelled by say index i , the intersection in eq. (3.1) is now trivial, containing only the identity. It can then be argued that the subspace, $\mathcal{H}_{\text{ginv}}^i$, of states in this sector admits a tensor product decomposition

$$\mathcal{H}_{\text{ginv}}^i = \mathcal{H}_{\text{ginv},\text{in}}^i \otimes \mathcal{H}_{\text{ginv},\text{out}}^i \quad (3.2)$$

with the operators in $\mathcal{A}_{\text{in}}, \mathcal{A}_{\text{out}}$ in the sector taking the form

$$\mathcal{O}_{\text{in}}^i \otimes \mathbf{I}_{\text{out}}^i, \mathbf{I}_{\text{in}}^i \otimes \mathcal{O}_{\text{out}}^i \quad (3.3)$$

respectively. The full Hilbert space can then be decomposed as

$$\mathcal{H}_{\text{ginv}} = \bigoplus_i \mathcal{H}_{\text{ginv}}^i = \bigoplus_i \mathcal{H}_{\text{ginv},\text{in}}^i \otimes \mathcal{H}_{\text{ginv},\text{out}}^i \quad (3.4)$$

and the density matrix of the inside ρ_{in} can be written as a sum over components in the different sectors,

$$\rho_{\text{in}} = \oplus_i \rho_{\text{in}}^i, \quad (3.5)$$

where each component ρ_{in}^i is obtained by starting with $\mathcal{H}_{\text{ginv}}^i$ and tracing over $\mathcal{H}_{\text{ginv},\text{out}}^i$.

Depending on which choice is made for the centre, one gets different definitions of the entanglement entropy. As mentioned above, the electric centre choice of CHR leads to a definition of entanglement entropy that coincides with the extended Hilbert space definition given above in the Abelian case. The index i introduced above for different sectors in this case coincides with the vector \mathbf{k} which gives the electric flux entering the inside at each boundary vertex.

From our discussion in section 2.1, see also GST, it follows that in the Abelian case by working in the extended Hilbert space \mathcal{H} and considering the space of gauge-invariant states $\mathcal{H}_{\text{ginv}} \subset \mathcal{H}$, we get

$$\mathcal{H}_{\text{ginv}} = \bigoplus_{\mathbf{k}} \mathcal{H}_{\text{ginv},\text{in}}^{\mathbf{k}} \otimes \mathcal{H}_{\text{ginv},\text{out}}^{\mathbf{k}}. \quad (3.6)$$

Here $\mathcal{H}_{\text{ginv},\text{in}}^{\mathbf{k}} \subset \mathcal{H}_{\text{in}}^{\mathbf{k}}$, consists of all states lying in $\mathcal{H}_{\text{in}}^{\mathbf{k}}$ which are invariant under gauge transformations acting on the inside vertices. Similarly, $\mathcal{H}_{\text{ginv},\text{out}}^{\mathbf{k}} \subset \mathcal{H}_{\text{out}}^{\mathbf{k}}$ consists of states invariant under gauge transformations acting on outside vertices.

3.2 Non-Abelian case

We now turn to examining the electric centre definition in the Non-Abelian case. For concreteness we work again with the SU(2) gauge theory; the generalisation to other Non-Abelian groups is quite straightforward. It is clear that here too there is a non-trivial centre in the algebra of gauge-invariant operators, $\mathcal{A}_{\text{in}}, \mathcal{A}_{\text{out}}$, which act on the inside and outside links. The choice for the centre we make, which is the analogue of the electric centre choice in the Abelian case, is given by $\{\hat{J}_i^2\}$, where this set contains the quadratic Casimirs at each boundary vertex V_i ³

$$\hat{J}_i^2 = (\hat{J}_{T,\text{in},i})^2. \quad (3.7)$$

Physically $\{\hat{J}_i^2\}$ is a gauge-invariant way to measure the electric flux entering a boundary vertex.

That these operators lie in the centre follows from noting that on gauge-invariant states, eq. (2.19) is valid and also that $(\hat{J}_{T,\text{out},i})^2, (\hat{J}_{T,\text{in},i})^2$ obviously commute with operators lying in $\mathcal{A}_{\text{in}}, \mathcal{A}_{\text{out}}$, respectively. Diagonalising the operators, eq. (3.7) and working in sectors consisting of eigenstates with fixed eigenvalues of the centre then leads to superselection sectors, specified by a vector \mathbf{k} , with k_i , eq. (2.21), specifying the eigenvalue of \hat{J}_i^2 at V_i . So far everything in the discussion is entirely analogous to what was done in the extended Hilbert space case; it also ties in with the general discussion of the CHR approach in the beginning of this section.

On general grounds, mentioned at the beginning of the section, one expects that the states of $\mathcal{H}_{\text{ginv}}$ lying in each of these sectors should now admit a tensor product decomposition of the form eq. (3.2), with the label i being identified with \mathbf{k} . The gauge-invariant operators restricted to this sector must also take the form, eq. (3.3).

To compare the electric centre definition we are developing here and the extended Hilbert space description above we now relate the spaces $\mathcal{H}_{\text{ginv},\text{in}}^{\mathbf{k}}, \mathcal{H}_{\text{ginv},\text{out}}^{\mathbf{k}}$, with those arising in the extended Hilbert space description discussed in section 2.2.

Notice that the extended Hilbert space \mathcal{H} can be written as

$$\mathcal{H} = \bigoplus \mathcal{H}^{\mathbf{k}}, \quad (3.8)$$

where

$$\mathcal{H}^{\mathbf{k}} = \mathcal{H}_{\text{in}}^{\mathbf{k}} \otimes \mathcal{H}_{\text{out}}^{\mathbf{k}}. \quad (3.9)$$

Just as $\mathcal{H}_{\text{in}}^{\mathbf{k}}$ can be expanded as eq. (2.22) so also $\mathcal{H}_{\text{out}}^{\mathbf{k}}$ can be written as,

$$\mathcal{H}_{\text{out}}^{\mathbf{k}} = \mathcal{H}'^1 \otimes \cdots \otimes \mathcal{H}'^i \otimes \cdots \otimes \mathcal{H}'^n \otimes \hat{\mathcal{H}}_{\text{out}}^{\mathbf{k}}. \quad (3.10)$$

The dimensions of \mathcal{H}^i , eq. (2.22) and \mathcal{H}'^i , eq. (3.10) are the same for $i = 1, \cdots, n$, and equal to $2j_i + 1$.

³This fact was also pointed out in [15].

There is a subspace $\hat{\mathcal{H}}_{\text{ginv},\text{in}}^{\mathbf{k}}$ of $\hat{\mathcal{H}}_{\text{in}}^{\mathbf{k}}$ which is gauge-invariant with respect to all gauge transformations acting on the inside vertices, and similarly $\hat{\mathcal{H}}_{\text{ginv},\text{out}}^{\mathbf{k}} \subset \hat{\mathcal{H}}_{\text{out}}^{\mathbf{k}}$ which is gauge-invariant with respect to all gauge transformations on the outside vertices.

We now argue that the spaces $\mathcal{H}_{\text{ginv},\text{in}}^{\mathbf{k}}, \mathcal{H}_{\text{ginv},\text{out}}^{\mathbf{k}}$, can be identified with $\hat{\mathcal{H}}_{\text{ginv},\text{in}}^{\mathbf{k}}, \hat{\mathcal{H}}_{\text{ginv},\text{out}}^{\mathbf{k}}$, respectively.

To show this we consider a gauge-invariant state $|\psi\rangle$ which lies in the sector \mathbf{k} . This state can be embedded in the $\mathcal{H}^{\mathbf{k}}$ sector of the extended Hilbert space. In fact, since the state is invariant with respect to gauge transformations acting on the inside and outside vertices it lies in the subspace

$$\mathcal{H}'^1 \otimes \cdots \otimes \mathcal{H}'^n \otimes \hat{\mathcal{H}}_{\text{ginv},\text{out}}^{\mathbf{k}} \otimes \mathcal{H}^1 \otimes \cdots \otimes \mathcal{H}^n \otimes \hat{\mathcal{H}}_{\text{ginv},\text{in}}^{\mathbf{k}}. \quad (3.11)$$

Gauge-invariance at the boundary vertices requires that the components in $\mathcal{H}^1, \mathcal{H}'^1$, and similarly for all the other boundary vertices, pair up to be singlets under the boundary gauge transformations. This means that any state $|\psi\rangle$ must be of the form

$$|\psi\rangle = \left(\sum_{\alpha_1, \alpha'_1} c_{\alpha_1, \alpha'_1} |\alpha_1\rangle \otimes |\alpha'_1\rangle \right) \otimes \left(\sum_{\alpha_2, \alpha'_2} c_{\alpha_2, \alpha'_2} |\alpha_2\rangle \otimes |\alpha'_2\rangle \right) \cdots \left(\sum_{\alpha_n, \alpha'_n} c_{\alpha_n, \alpha'_n} |\alpha_n\rangle \otimes |\alpha'_n\rangle \right) \\ \otimes \sum_{A, b} C_{Ab} |A\rangle_{\text{in}} \otimes |b\rangle_{\text{out}}. \quad (3.12)$$

Here $|\alpha_1\rangle, |\alpha'_1\rangle$, with $\alpha_1, \alpha'_1 = 1, \dots, 2j_1 + 1$ are a basis for $\mathcal{H}^1, \mathcal{H}'^1$ respectively and so on. The coefficients c_{α_1, α'_1} are such that the state is a singlet with respect to gauge transformations at the vertex V_1 and similarly for the other boundary vertices. In the last term $\{|A\rangle\}, \{|b\rangle\}$ are bases in $\hat{\mathcal{H}}_{\text{ginv},\text{in}}^{\mathbf{k}}, \hat{\mathcal{H}}_{\text{ginv},\text{out}}^{\mathbf{k}}$, respectively, so this term gives a state in $\hat{\mathcal{H}}_{\text{ginv},\text{in}}^{\mathbf{k}} \otimes \hat{\mathcal{H}}_{\text{ginv},\text{out}}^{\mathbf{k}}$. Note that for all states in this sector the first n terms in the tensor product above remain the same and only the last term which involves the state in $\hat{\mathcal{H}}_{\text{ginv},\text{in}}^{\mathbf{k}} \otimes \hat{\mathcal{H}}_{\text{ginv},\text{out}}^{\mathbf{k}}$ changes.

This shows that there is a one to one correspondence between states in the space $\mathcal{H}_{\text{ginv}}^{\mathbf{k}}$, which arose in the electric centre discussion and states in $\hat{\mathcal{H}}_{\text{ginv},\text{in}}^{\mathbf{k}} \otimes \hat{\mathcal{H}}_{\text{ginv},\text{out}}^{\mathbf{k}}$, which we defined using the extended Hilbert space. The map is one-to-one because given any state in $\hat{\mathcal{H}}_{\text{ginv},\text{in}}^{\mathbf{k}} \otimes \hat{\mathcal{H}}_{\text{ginv},\text{out}}^{\mathbf{k}}$, we can obtain a state in $\mathcal{H}_{\text{ginv}}^{\mathbf{k}}$ by appending the first n terms which ensure gauge-invariance with respect to boundary gauge transformations. Moreover from the form of a gauge-invariant operator $\mathcal{O}_{\text{in}}^{\mathbf{k}}$, eq. (2.24) which acts on inside links and a similar form for operators acting on the outside links we see that acting on $\hat{\mathcal{H}}_{\text{ginv},\text{in}}^{\mathbf{k}} \otimes \hat{\mathcal{H}}_{\text{ginv},\text{out}}^{\mathbf{k}}$ these operators take the form, $\hat{\mathcal{O}}_{\text{in}}^{\mathbf{k}} \otimes \mathbf{I}_{\text{out}}, \mathbf{I}_{\text{in}} \otimes \hat{\mathcal{O}}_{\text{out}}^{\mathbf{k}}$ respectively in $\hat{\mathcal{H}}_{\text{ginv},\text{in}}^{\mathbf{k}} \otimes \hat{\mathcal{H}}_{\text{ginv},\text{out}}^{\mathbf{k}}$.

It then follows, from the general arguments at the beginning of the section, that $\mathcal{H}_{\text{ginv},\text{in}}^{\mathbf{k}}, \mathcal{H}_{\text{ginv},\text{out}}^{\mathbf{k}}$, which we defined in the electric centre discussion above can be identified with $\hat{\mathcal{H}}_{\text{ginv},\text{in}}^{\mathbf{k}}, \hat{\mathcal{H}}_{\text{ginv},\text{out}}^{\mathbf{k}}$, respectively,

$$\mathcal{H}_{\text{ginv},\text{in}}^{\mathbf{k}} \simeq \hat{\mathcal{H}}_{\text{ginv},\text{in}}^{\mathbf{k}}, \quad (3.13)$$

$$\mathcal{H}_{\text{ginv},\text{out}}^{\mathbf{k}} \simeq \hat{\mathcal{H}}_{\text{ginv},\text{out}}^{\mathbf{k}}. \quad (3.14)$$

We remind the reader that $\hat{\mathcal{H}}_{\text{ginv},\text{in}}^{\mathbf{k}}, \hat{\mathcal{H}}_{\text{ginv},\text{out}}^{\mathbf{k}}$ were defined by working in the extended Hilbert space, \mathcal{H} .

As a result, it follows from eq. (3.4) and eq. (3.13), eq. (3.14), that the Hilbert space of gauge-invariant states can be expressed as a sum

$$\mathcal{H}_{\text{ginv}} = \bigoplus \hat{\mathcal{H}}_{\text{ginv},\text{in}}^{\mathbf{k}} \otimes \hat{\mathcal{H}}_{\text{ginv},\text{out}}^{\mathbf{k}}. \quad (3.15)$$

3.3 Entanglement entropy

We now come to the main point of this section, namely the difference between the entanglement entropy which arises from the electric centre description given above and the extended Hilbert space definition of the entanglement entropy.

To calculate the difference we need to relate the density matrix ρ_{in} in the two cases. The density matrix in the superselection sector \mathbf{k} for the extended Hilbert space is given in eq. (2.23). For a gauge-invariant state $|\psi\rangle$, it is easy to see that $\hat{\rho}_{\text{in}}^{\mathbf{k}}$ has support only on $\hat{\mathcal{H}}_{\text{ginv},\text{in}}^{\mathbf{k}} \subset \hat{\mathcal{H}}_{\text{in}}^{\mathbf{k}}$. Up to a normalisation, we now argue that the density matrix in a sector \mathbf{k} in the electric centre definition is in fact $\hat{\rho}_{\text{in}}^{\mathbf{k}}$ restricted to $\hat{\mathcal{H}}_{\text{ginv},\text{in}}^{\mathbf{k}}$. By construction the expectation value for any gauge-invariant operator can be calculated by working in the extended Hilbert space. For an operator acting on the inside links we get

$$\langle \mathcal{O}_{\text{in}} \rangle = \sum_{\mathbf{k}} \text{Tr}_{\mathcal{H}_{\text{in}}^{\mathbf{k}}} \mathcal{O}_{\text{in}}^{\mathbf{k}} \rho_{\text{in}}^{\mathbf{k}}. \quad (3.16)$$

From eq. (2.23), eq. (2.24), we see that

$$\text{Tr}_{\mathcal{H}_{\text{in}}^{\mathbf{k}}} \mathcal{O}_{\text{in}}^{\mathbf{k}} \rho_{\text{in}}^{\mathbf{k}} = \left(\prod_i (2j_i + 1) \right) \text{Tr}_{\hat{\mathcal{H}}_{\text{ginv},\text{in}}^{\mathbf{k}}} \hat{\mathcal{O}}_{\text{in}}^{\mathbf{k}} \hat{\rho}_{\text{in}}^{\mathbf{k}}, \quad (3.17)$$

where we have used the fact that the trace on the r.h.s. can be restricted to $\hat{\mathcal{H}}_{\text{ginv},\text{in}}^{\mathbf{k}}$, since \mathcal{O} is gauge-invariant and $\rho_{\text{in}}^{\mathbf{k}}$ has support only in $\hat{\mathcal{H}}_{\text{ginv},\text{in}}^{\mathbf{k}}$. From eq. (2.8), eq. (2.9), eq. (2.27) it then follows that the density matrix in sector \mathbf{k} in the electric centre definition is given by

$$\rho_{\text{in},\text{ec}}^{\mathbf{k}} = p_{\mathbf{k}} \bar{\rho}_{\text{in}}^{\mathbf{k}} \quad (3.18)$$

From eq. (2.28) and the fact that $\bar{\rho}_{\text{in}}^{\mathbf{k}}$ has support on only $\hat{\mathcal{H}}_{\text{ginv},\text{in}}^{\mathbf{k}}$ it follows that

$$\text{Tr}_{\hat{\mathcal{H}}_{\text{ginv},\text{in}}^{\mathbf{k}}} \bar{\rho}_{\text{in}}^{\mathbf{k}} = 1 \quad (3.19)$$

It then follows that the entanglement entropy which in the electric centre definition is given by

$$S_{EE,\text{ec}} = - \sum_{\mathbf{k}} \text{Tr}_{\hat{\mathcal{H}}_{\text{ginv},\text{in}}^{\mathbf{k}}} \rho_{\text{in},\text{ec}}^{\mathbf{k}} \log \rho_{\text{in},\text{ec}}^{\mathbf{k}}, \quad (3.20)$$

which becomes

$$S_{EE,\text{ec}} = - \sum_{\mathbf{k}} p_{\mathbf{k}} \log p_{\mathbf{k}} - \sum_{\mathbf{k}} p_{\mathbf{k}} \text{Tr}_{\hat{\mathcal{H}}_{\text{ginv},\text{in}}^{\mathbf{k}}} \bar{\rho}_{\text{in}}^{\mathbf{k}} \log \bar{\rho}_{\text{in}}^{\mathbf{k}}. \quad (3.21)$$

Comparing we see that the second term in eq. (2.29) is missing. More generally, the second term in eq. (2.30) will be missing, as was mentioned at the start of this section.⁴ The physical origin of this missing term was discussed in section 2.2. It arises in the extended Hilbert space definition because the total angular momentum carried by inside and outside links meeting at a boundary vertex must sum to zero, giving rise to non-zero entanglement between the inside and outside links. In the electric centre definition we work with the Hilbert space of gauge-invariant states $\mathcal{H}_{\text{ginv}}$ which can be decomposed only in terms of $\hat{\mathcal{H}}_{\text{ginv},\text{in}}^k, \hat{\mathcal{H}}_{\text{ginv},\text{out}}^k$, eq. (3.15). As a result this additional contribution is absent.

4 Entanglement distillation and dilution

It is a standard practice to compare the entanglement in a bipartite system with a reference system typically taken to be a set of Bell pairs. The comparison is done using the processes of entanglement dilution and distillation which involve only Local Operations and Classical Communication (LOCC), see [16–19]. For a system with local degrees of freedom, like a spin system, where the Hilbert space of states admits a tensor product decomposition, it is well known that the number of Bell pairs which are consumed in dilution or produced in distillation, is given by the entanglement entropy,

$$S_{EE} = -\text{Tr}(\rho \log(\rho)). \quad (4.1)$$

More accurately, this statement is true in the asymptotic limit where we take n copies of the system and find that the number of pairs used up in dilution, k' , and those produced in distillation, k , both tend to the limit,

$$\lim_{n \rightarrow \infty} \frac{k'}{n} = \lim_{n \rightarrow \infty} \frac{k}{n} = S_{EE}. \quad (4.2)$$

where k' approaches the limit from above and k from below. The starting point for distillation is that Alice and Bob share n copies of the state $|\psi\rangle$ and each has k unentangled reference qubits. Distillation is the process of converting the qubits into k Bell pairs by using up the entanglement in $|\psi\rangle$. The set-up for dilution is that Alice and Bob share k' Bell pairs and Alice has n copies of the bipartite system AC in the state $|\psi\rangle$ at her disposal. By LOCC operations the state of the system C is then teleported to Bob, using the Bell pairs, resulting in the n copies of state $|\psi\rangle$ now being shared between Alice and Bob. The minimum number of Bell pairs required for this is k' . We then find that eq. (4.2) holds (these statements can be made more precise using ϵ 's and δ 's and the analysis below will be done in this more careful manner).

In contrast, it was argued in CHR and GST that for the extended Hilbert space definition of entanglement entropy eq. (4.2) is no longer true for gauge theories. This is because of the presence of superselection sectors which cannot be changed by local gauge-invariant operations, as reviewed in section 2. The existence of superselection sectors imposes an essential limitation on extracting the full entanglement entropy using dilution

⁴The trace in the last term in eq. (2.29), eq. (2.30) can be restricted to $\hat{\mathcal{H}}_{\text{ginv},\text{in}}^k$ since $\hat{\rho}_{\text{in}}^k$ has support only on this subspace of $\hat{\mathcal{H}}_{\text{in}}^k$.

or distillation. In the Abelian case the entanglement entropy can be written as a sum of two terms given in eq. (2.11). Similarly in the Non-Abelian case it can be written as the sum of three terms given in eq. (2.30). One expects that the first term in eq. (2.11) in the Abelian case, and the first two terms in eq. (2.30) for the Non-Abelian case cannot be extracted using local gauge-invariant operations.

In this section we will carry out a more detailed investigation and find that this expectation is indeed borne out. For distillation we will find that the state shared between Alice and Bob at the end is still partially entangled with a residual entanglement entropy that cannot be reduced any further. This entropy is indeed given by the first term in eq. (2.11) for the Abelian case and the sum of the first and second terms of eq. (2.30) in the Non-Abelian case. In dilution we will find that Alice and Bob need to start with a partially entangled state so that at the end of the dilution process they can share n copies of the state $|\psi\rangle$. The entanglement in this starting state in dilution, per copy, will be the same as the residual entanglement left after distillation. The number of Bell pairs extracted in distillation and consumed in dilution in this way will turn out to be equal asymptotically,

$$\lim_{N \rightarrow \infty} \frac{k'}{N} = \lim_{N \rightarrow \infty} \frac{k}{N} \quad (4.3)$$

But this limit will not equal S_{EE} . Instead, the limit, i.e., maximum number of Bell pairs which can be extracted, will equal the second term in eq. (2.11) for the Abelian case, and the third term in eq. (2.30) in the Non-Abelian case.

A similar question relating entanglement to dilution and distillation for a simpler superselection rule, total number conservation, was answered by Schuch, Verstraete and Cirac (SVC) in [20, 21]. In that case, local operations cannot change the total number of particles on either Alice or Bob's side. If a state has 1 particle on each side, then n copies of the state can be transformed to another state with n particles on each side; that is, there is still just one superselection rule. In our case, however, each copy has superselection sectors arising from the fact that the Gauss law constraint must be met in every copy. As a result, with n copies we get n sets of superselection sectors, one for each copy.

4.1 A toy model

We will find it convenient to carry out the analysis in a toy model consisting of just 4 qubits. It will become clear as we proceed that our conclusion can be easily generalised also to Abelian and Non-Abelian gauge theories. In the discussion below, at the risk of introducing somewhat confusing conventions, we label a basis for each qubit, as $|1\rangle$ and $|0\rangle$, with $\sigma^3 |1\rangle = |1\rangle, \sigma^3 |0\rangle = -|0\rangle$.

We will regard this 4 qubit system, as a bipartite system, with the first two qubits being the first part, A , and the third and fourth qubits being the second part, B . In order to model the constraint of gauge-invariance we will impose a condition on this system. Namely, that the only allowed states are those for which the second and third qubits take the same value. Thus an allowed or physical state in our toy model has the form

$$|\psi\rangle = \sum_{i,j,k=0}^1 \sqrt{p_j} a_{ijk} |i, j\rangle_A |j, k\rangle_B = \sum_j \sqrt{p_j} |\psi_j\rangle_{AB}, \quad (4.4)$$

where

$$|\psi_j\rangle_{AB} = \sum_{i,k=0}^1 a_{ijk} |i, j\rangle_A |j, k\rangle_B \quad (4.5)$$

and we have normalised $|\psi_j\rangle$ so that

$$\langle\psi_j|\psi_j\rangle = 1. \quad (4.6)$$

The constraint that the middle two qubits in eq. (4.4) are the same is the analogue of Gauss' law, which in a gauge theory without matter implies that the electric flux leaving a region is the same as that entering its complement.

The allowed operators in this toy model are those which preserve the constraint and therefore change the middle two qubits together so that they continue to be equal. This is analogous the fact that the allowed operators in a gauge theory are gauge-invariant. As a result, the full algebra of allowed operators is generated by

$$\mathcal{G} = \{\sigma_1^z, \sigma_1^x, \sigma_2^z, \sigma_2^x \otimes \sigma_3^x, \sigma_3^z, \sigma_4^z, \sigma_4^x\}. \quad (4.7)$$

Here we are using the notation that the superscript denotes the kind of Pauli matrices, $\sigma^x, \sigma^y, \sigma^z$, while the subscript indicates which qubit the matrices act on.

There are two subalgebras, \mathcal{A} and \mathcal{B} , which act on A and B , of allowed sets of operators, generated by

$$\mathcal{G}_A = \{\sigma_1^z, \sigma_1^x, \sigma_2^z\}, \quad (4.8)$$

and

$$\mathcal{G}_B = \{\sigma_3^z, \sigma_4^z, \sigma_4^x\}, \quad (4.9)$$

respectively. The centre of each subalgebra is generated by $\{\sigma_2^z = \sigma_3^z\}$. The operators in \mathcal{A} and \mathcal{B} are the analogue of gauge-invariant operators which act entirely in the inside or outside links in the gauge theory. The operator $\sigma_2^x \otimes \sigma_3^x$ is the analogue of a Wilson loop which crosses from one region to its complement changing the electric flux.

In the discussion below, in an abuse of terminology, to emphasise the analogy we will sometimes refer to the allowed operators in the algebra generated by \mathcal{G} as gauge-invariant operators and to the operators in the subalgebras \mathcal{A}, \mathcal{B} as local gauge-invariant operators.

There are two superselection sectors in this toy model given by two values of the middle two qubits, $j = 1, 0$ in eq. (4.4). It is clear that the subalgebras \mathcal{A}, \mathcal{B} , acting on A or B cannot change these superselection sectors. An important comment worth noting is that if we consider n copies of $|\psi\rangle$, each copy will have its own superselection sectors, resulting in a total of 2^n superselection sectors.

Let us end with a few more important comments. Starting from $|\psi\rangle$ in eq. (4.4) and tracing over B we get a density matrix

$$\rho = \begin{bmatrix} \rho_0 & 0 \\ 0 & \rho_1 \end{bmatrix} \quad (4.10)$$

for the state in A . ρ is block diagonal with the two blocks referring to the two superselection sectors where the second qubit takes values 0, 1 respectively. Each block is a 2×2 matrix. ρ_0, ρ_1 are the unnormalised density matrices in these two sectors with

$$\text{Tr}(\rho_0) = p_0, \quad (4.11)$$

$$\text{Tr}(\rho_1) = p_1, \quad (4.12)$$

being the probabilities for the two superselection sectors.

Now consider an allowed local unitary operator \hat{U} . Under the action of this operator

$$\rho \rightarrow \hat{U} \rho \hat{U}^\dagger. \quad (4.13)$$

Since U cannot change the superselection sector it is therefore also block diagonal. We then conclude that any such operator leaves p_0, p_1 unchanged. Similarly, consider a generalised measurement. This corresponds to a set of operators, $\{M_i\}, i = 1, \dots, N$ with

$$\sum_i M_i^\dagger M_i = \mathbf{1}. \quad (4.14)$$

Under it the density matrix transforms as

$$\rho \rightarrow \sum_i M_i \rho M_i^\dagger. \quad (4.15)$$

An allowed measurement must be block diagonal in the basis above since it cannot change the superselection sector (i.e. each M_i must be block diagonal). It then follows from eq. (4.15) that this also leaves p_0, p_1 unchanged, as discussed in SVC.

Next, let us write the state $|\psi\rangle$, eq. (4.4) in its Schmidt basis in every sector,

$$|\psi_j\rangle = \sum_{I=0}^1 \sqrt{\alpha_{I,j}} |I^j, j\rangle_A |j, I^j\rangle_B, \quad (4.16)$$

Note that the I label takes two values, and has a superscript when it appears as a label for states because the Schmidt basis for the first and fourth qubit is independent in every superselection sector. E.g., $|0^0\rangle \neq |0^1\rangle$.

It is easy to see that the Von Neumann entropy of the density matrix for A in this state is

$$S = S_c + S_q, \quad (4.17)$$

where the first term is,

$$S_c = -p_0 \log p_0 - p_1 \log p_1, \quad (4.18)$$

and the second term is

$$S_q = -[p_0 \{\alpha_{1,0} \log \alpha_{1,0} + \alpha_{0,0} \log \alpha_{0,0}\}] - [p_1 \{\alpha_{1,1} \log \alpha_{1,1} + \alpha_{0,1} \log \alpha_{0,1}\}]. \quad (4.19)$$

These terms have the following interpretation. S_c is the entropy which arises due to the probability of being in different superselection sectors. In fact, it is the minimum

entropy which can arise from any density matrix, eq. (4.10), subject to the constraint eq. (4.11), eq. (4.12), which we have seen is preserved by any allowed unitary operator or measurement. S_q on the other hand is the average of the entanglement entropies in the two superselection sectors weighted by the probability of being in these sectors.

The distillation and dilution protocols we will discuss next involve local gauge-invariant operators. Since, as noted above, these do not change p_0, p_1 , we do not expect these protocols to be able to extract the entropy in the first term, S_c . Instead, an efficient protocol should at most be able to extract the second term, S_q . We will see that the protocols we discuss below indeed meet this expectation.

We had mentioned towards the end of section 2 that there are classical type of contributions which arise in the entanglement of gauge theories, corresponding to the first term in eq. (2.11) and the first two terms in eq. (2.30). The S_c term, eq. (4.18), in the toy model we are considering here is in fact analogous to those contributions⁵ eq. (2.30). We will return to the connection with gauge theories again at the end of this section after analysing the toy model in more detail.

4.2 The distillation protocol

We start by taking n copies of the state denoted by $|\psi\rangle^{\otimes n}$ with Alice having access to the n copies in A and Bob to the n copies in B . In the discussion below the different copies will be labelled by the index $r = 1, 2, \dots, n$. Next, we consider a fuzzy measurement that Alice performs on her n copies of the A subsystem, corresponding to the operator

$$\hat{O}_1 = \sum_{r=1}^n \left(\frac{1 + \sigma_2^{z,r}}{2} \right). \quad (4.20)$$

Here, we remind the reader that the label z denotes the σ^z Pauli matrix, the subscript 2 indicates that it acts on the 2nd qubit in eq. (4.4) and the label r specifies which copy among the n the operator acts on. If the result of this measurement is m_1 , then the state after the measurement is given by the symmetric superposition,

$$|\psi_1\rangle = \frac{1}{\sqrt{N_1}} \sum_{(j_r)|m_1} |\psi_{j_r}\rangle_A \otimes |\psi_{j_r}\rangle_B, \quad (4.21)$$

where $\sum_{(j_r)|m_1}$ indicates that the sum is over all lists (j_r) , subject to the constraint that $\sum_r j_r = m_1$. The normalisation N_1 in eq. (4.21) takes the value

$$N_1 = \binom{N}{m_1} \quad (4.22)$$

and is the total number of terms in the symmetric sum.

Next, Alice measures two additional variables,

$$\hat{O}_2 = \sum_{r=1}^n (|1^1\rangle_r \langle 1^1|_r) \left(\frac{1 + \sigma_2^{z,r}}{2} \right) \quad (4.23)$$

⁵Actually, as we will see later, for the Non-Abelian case S_c is analogous to the first term in eq. (2.30) which is also the classical type of contribution we get in the electric centre definition, eq. (3.21).

and

$$\hat{O}_3 = \sum_{r=1}^n (|1^0\rangle_r \langle 1^0|_r) \left(\frac{1 - \sigma_2^{z,r}}{2} \right). \quad (4.24)$$

Here $|1^1\rangle_r \langle 1^1|_r$ stands for a projector which acts on the first qubit of the r^{th} copy, with $|1^1\rangle$ denoting the state of the first qubit in the Schmidt basis for the superselection sector with $j = 1$, see discussion after eq. (4.16) above. Similarly $|1^0\rangle_r \langle 1^0|_r$ also stands for a projector of the r^{th} copy, this time to a Schmidt basis state in the $j = 0$ superselection sector

If these two measurements give rise to the results s_1 and t_1 respectively then the state after these measurements is a symmetric superposition of N_{T_1} terms

$$|\psi'\rangle = \frac{1}{\sqrt{N_{T_1}}} \sum_{(j_r, I_r)} \prod_r |I_r^{j_r}, j_r\rangle_A \otimes |j_r, I_r^{j_r}\rangle_B. \quad (4.25)$$

where the sum is over all lists (j_r, I_r) with the following properties:

$$\sum j_r = m_1 \quad (4.26)$$

$$\sum' I_r = s_1 \quad (4.27)$$

$$\sum'' I_r = t_1. \quad (4.28)$$

Here \sum' indicates a restricted sum over r where only those terms with $j_r = 1$ contribute, while \sum'' indicates a restricted sum where only those values of r with $j_r = 0$ contribute. Note that N_{T_1} in eq. (4.25) is given by

$$N_{T_1} = \binom{n}{m_1} \binom{m_1}{s_1} \binom{n - m_1}{t_1}. \quad (4.29)$$

Now we come to an important point. The N_T terms in eq. (4.25) arise as follows: there are $\binom{n}{m}$ distinct values that the set $\{j_1, j_2, \dots, j_n\}$ can take subject to the constraint that $\sum_r j_r = m$. Each of these choices corresponds to a different superselection sector. Working with only local gauge-invariant operators we do not expect to extract the entanglement between the qubits Alice and Bob share due to this choice of superselection sectors. For each choice of $\{j_1, j_2, \dots, j_n\}$ and each possible value of s and t we have $\binom{m}{s} \binom{n-m}{t}$ different states corresponding to different choices of the I labels. We would expect to be able to extract the entanglement in these states while preserving the superselection sectors for each copy. Below we see in a precise sense that this expectation is indeed met.

To begin, consider the case where

$$2^p = \binom{m_1}{s_1} \binom{n - m_1}{t_1}, \quad (4.30)$$

for some integer p . In this case we provide both Alice and Bob with p qubits which are unentangled and in the state

$$|0\rangle_A^{\otimes p} \otimes |0\rangle_B^{\otimes p}, \quad (4.31)$$

where A and B refer to the p qubits in Alice and Bob's possession respectively. A unitary transformation can now be carried out in each superselection sector, i.e. in each sector

labelled by a distinct choice of the set $\{j_1, j_2 \dots j_n\}$, by Alice which swaps the component of $|\psi'\rangle$, eq. (4.25), in this superselection sector, which she has access to, with that of the p qubits. Bob carries out a similar swap on his side as well. Since the state in eq. (4.25) is a symmetric superposition with the same number of summands, 2^p , in each superselection sector this results in maximally entangling the two sets of p qubits. The resulting state of the full system consisting of the original n copies and the additional qubits is then given by

$$\left(\frac{1}{\sqrt{N_1}} \sum_{(j_r)|m_1} \prod_r |0^{j_r}, j_r\rangle_A \otimes |j_r, 0^{j_r}\rangle_B \right) \otimes |\Phi_+\rangle^{\otimes p}, \quad (4.32)$$

where

$$|\Phi_+\rangle = (|00\rangle + |11\rangle)/\sqrt{2} \quad (4.33)$$

refers to a Bell pair shared between Alice and Bob.

We emphasise that the components of $|\psi'\rangle$, eq. (4.25), corresponding to different superselection sectors, have the same number of terms in a symmetric superposition. This ensures that the above swap leaves the reference qubits unentangled with the state of the system.

Let us also note that a swap transformation, which we will also often use below, is a unitary transformation defined as follows. Let \mathcal{H} and $\tilde{\mathcal{H}}$ be two Hilbert spaces of the same dimension and $|\phi\rangle \otimes |\chi\rangle \in \mathcal{H} \otimes \tilde{\mathcal{H}}$, then under a swap transformation S which takes $\mathcal{H} \otimes \tilde{\mathcal{H}} \rightarrow \mathcal{H} \otimes \tilde{\mathcal{H}}$,

$$|\phi\rangle \otimes |\chi\rangle \rightarrow |\chi\rangle \otimes |\phi\rangle. \quad (4.34)$$

In the limit when we start with a large number of copies, $n \gg 1$, it is easy to see that with probability $1 - \epsilon$

$$2^{nS_q(1-\delta)} < \binom{m_1}{s_1} \binom{n-m_1}{t_1} < 2^{nS_q(1+\delta)}, \quad (4.35)$$

where S_q is the entropy eq. (4.19), and δ is small,

$$\delta \sim \frac{1}{\sqrt{n}}. \quad (4.36)$$

We also remind the reader that in eq. (4.19) $\alpha_{I,j}$ are the coefficients which arise in the expansion in the Schmidt basis, eq. (4.16), and p_0, p_1 are the probabilities for the superselection sectors $j = 0, 1$ respectively, eq. (4.4).

Eq. (4.35) arises by noting that for $n \gg 1$ the expectation value for m_1 is

$$\langle m_1 \rangle = np_1 \quad (4.37)$$

and for s_1, t_1 are

$$\langle s_1 \rangle = np_1 \alpha_{1,1} \quad (4.38)$$

and

$$\langle t_1 \rangle = np_0 \alpha_{1,0} \quad (4.39)$$

respectively. And by noting that fluctuations about this mean are suppressed by a fractional power of order $1/\sqrt{n}$. We also note that p defined in eq. (4.30) satisfies the condition

$$p > nS_q(1 - \delta), \quad (4.40)$$

which follows from eq. (4.35).

Of course, in general p defined by eq. (4.30) will not be an integer. In this case we can repeat this whole procedure multiple times. Each time starting with n copies of the state $|\psi\rangle$ Alice measures operators $\hat{O}_1, \hat{O}_2, \hat{O}_3$, obtaining in the l^{th} iteration values m_l, s_l, t_l respectively. After L total iterations the total number of copies is

$$n_T = nL \quad (4.41)$$

and the resulting state will consist of a symmetric superposition of several superselection sectors. The number of terms in each superselection sector now will be

$$N_T = \binom{m_1}{s_1} \binom{n-m_1}{t_1} \binom{m_2}{s_2} \binom{n-m_2}{t_2} \cdots \binom{m_L}{s_L} \binom{n-m_L}{t_L}. \quad (4.42)$$

For L sufficiently large one can in this way ensure that

$$2^k \leq N_T \leq 2^k(1 + \epsilon) \quad (4.43)$$

for some integer k . Since $2^k(1 + \epsilon) < 2^{k+1}$, to proceed Alice and Bob can then take $k+1$ reference qubits each, which are initially totally unentangled in a state analogous to eq. (4.31), and swap their state with that of the system in any given superselection sector. More accurately, Alice carries out this swap between the reference qubits and the component of the state of the system she has access to and similarly for Bob. A measurement can then be made projecting the state of the qubits to a 2^k dimensional subspace with probability $1 - \epsilon$. This will lead to k extracted Bell pairs which are in a totally entangled state.

It follows from eq. (4.35), eq. (4.19) that the ratio

$$\lim_{n_T \rightarrow \infty} \frac{k}{n_T} = S_q \quad (4.44)$$

This distillation procedure will work with probability $1 - \epsilon$, as mentioned above. With a large enough number of copies, ϵ can be made arbitrarily small.

Note that in eq. (4.32) the two parts of the n copies of the original system still share some residual entanglement, since the choice of superselection sectors in both parts is correlated. This is also true more generally if we repeat the procedure L times. The residual entanglement is a consequence of the superselection sectors, i.e. of gauge-invariance. Local gauge-invariant operations cannot change the superselection sectors and thus acting with only these operators does not allow this remaining entanglement to be extracted. The final state of the original system, after L iterations is

$$|\psi^{\text{fin}}\rangle = \left(\frac{1}{\sqrt{N_1}} \sum_{(j_{r_1})|m_1} \prod_{r_1} |0^{j_{r_1}}, j_{r_1}\rangle_A \otimes |j_{r_1}, 0^{j_{r_1}}\rangle_B \right) \otimes \cdots \otimes \left(\frac{1}{\sqrt{N_l}} \sum_{(j_{r_l})|m_l} \prod_{r_l} |0^{j_{r_l}}, j_{r_l}\rangle_A \otimes |j_{r_l}, 0^{j_{r_l}}\rangle_B \right) \\ \otimes \cdots \otimes \left(\frac{1}{\sqrt{N_L}} \sum_{(j_{r_L})|m_L} \prod_{r_L} |0^{j_{r_L}}, j_{r_L}\rangle_A \otimes |j_{r_L}, 0^{j_{r_L}}\rangle_B \right), \quad (4.45)$$

where

$$N_l = \binom{n}{m_l}. \quad (4.46)$$

4.3 The dilution protocol

In the usual discussion of dilution one considers a situation where at the start Alice and Bob share a large number of Bell pairs and also Alice has at her disposal n copies of the state $|\psi\rangle$. The state $|\psi\rangle$ is a bipartite system whose two parts, both in possession of Alice, we refer to as A' and C . Local operations are now carried out on the n copies of C by Alice and local operations are also carried out by Bob in a coordinated way made possible by classical communication. We then ask what is the minimum number of Bell pairs that are needed so that after the protocol is over the entanglement between A' and C is now shared between Alice and Bob. More precisely, what is the minimum number of Bell pairs needed, as $n \rightarrow \infty$, so that Alice and Bob share the n copies of $|\psi\rangle$ with very high fidelity.

The new feature in our discussion is that we are dealing with a system with superselection sectors. Local allowed operators cannot change these superselection sectors. Thus acting with only these operators we cannot hope to extract all the entanglement entropy eq. (4.17) between A' and C and transfer it to lie between Alice and Bob.

In order to construct the state $|\psi\rangle$ with high fidelity shared between Alice and Bob it will therefore be necessary to start with a partially entangled state between Alice and Bob. This state will carry the “barebones” entanglement given by the first term eq. (4.18) which we do not expect to be able to transfer. Then we will show that the rest of the entanglement, eq. (4.19), in the asymptotic limit of a large number of copies, n , with high fidelity can be obtained by dilution of a set of shared Bell pairs. It will turn out that the minimum number of Bell pairs required in this process k' indeed meets the condition

$$\lim_{n \rightarrow \infty} \frac{k'}{n} = \lim_{n_T \rightarrow \infty} \frac{k}{n_T} = S_q, \quad (4.47)$$

where k, n_T are defined near eq. (4.44) and S_q is defined in eq. (4.19).

The partially entangled state we will start with in the protocol is given by

$$|\psi^{\text{in}}\rangle = \frac{1}{\sqrt{D}} \sum_{m=\bar{m}(1-\delta)}^{m=\bar{m}(1+\delta)} \sum_{(j_r)|m} \prod_r \sqrt{p_r} |0^{j_r}, j_r\rangle_A \otimes |j_r, 0^{j_r}\rangle_B. \quad (4.48)$$

Here, D is the overall normalisation required to make this state of unit norm,

$$\bar{m} = np_1 \quad (4.49)$$

is the expectation value of $\sum j_r$ in the $n \rightarrow \infty$ limit, and

$$\delta \sim O(1/\sqrt{n}). \quad (4.50)$$

The state in eq. (4.48) is a bipartite state whose two parts denoted by A and B are accessible to Alice and Bob respectively.

The rest of the steps involved in the protocol are similar to the standard discussion of dilution. We start with n copies of $|\psi\rangle$ whose two parts we refer to as A' and C both of which are in possession of Alice. From eq. (4.4) we see that

$$|\psi\rangle^{\otimes n} = \sum_{(j_r)} \sum_{(I_r)} \prod_r \sqrt{p_r} \sqrt{\alpha_{I_r, j_r}} |I_r^{j_r}, j_r\rangle_{A'} \otimes |j_r, I_r^{j_r}\rangle_C. \quad (4.51)$$

Here $\sum_{(j_r)}$ stands for the sum over all lists $\{j_1, j_2, \dots, j_n\}$, and $\sum_{(I_r)}$ for the sum over all lists $\{I_1, I_2, \dots, I_n\}$.

Next, Alice Schumacher compresses the state $|\psi\rangle^{\otimes n}$ by projecting onto a typical subspace. This can be done by measuring suitable fuzzy operators. The state is then give by

$$|\Phi\rangle = \mathcal{N} \sum_{m=\bar{m}(1-\delta)}^{m=\bar{m}(1+\delta)} \sqrt{p_1^m} \sqrt{p_0^{n-m}} \sum_{s=\bar{s}(1-\delta_2)}^{s=\bar{s}(1+\delta_2)} \sum_{t=\bar{t}(1-\delta_3)}^{t=\bar{t}(1+\delta_3)} \sqrt{\alpha_{1,1}^s} \sqrt{\alpha_{0,1}^{m-s}} \sqrt{\alpha_{1,0}^t} \sqrt{\alpha_{0,0}^{n-m-t}} \sum_{(j_r, I_r)} \prod_r |I_r^{j_r}, j_r\rangle_{A'} \otimes |j_r, I_r^{j_r}\rangle_C, \quad (4.52)$$

where \mathcal{N} is an overall normalisation to make $|\Phi\rangle$ of unit norm, and \bar{m} and δ are given in eq. (4.49) and eq. (4.50) respectively. Also note that \bar{s}, \bar{t} are dependent on m and given by

$$\bar{s} = m\alpha_{1,1} \quad (4.53)$$

$$\bar{t} = (n-m)\alpha_{1,0}. \quad (4.54)$$

They are the expectation values for the operator \hat{O}_2, \hat{O}_3 , eq. (4.23), eq. (4.24), respectively for

$$\sum_r j_r = m. \quad (4.55)$$

Note also that we take $\delta_2, \delta_3 \sim O(1/\sqrt{n})$ in eq. (4.52). And finally that $\sum_{(j_r, I_r)}$ denotes a sum over lists $\{(j_1, I_1), (j_2, I_2), \dots, (j_n, I_n)\}$ such that $\sum_r j_r = m$; $\sum' I_r = s$, and $\sum'' I_r = t$, where \sum' denotes a sum in which only terms with $j_r = 1$ contributes, and \sum'' denotes a sum in which only terms with $j_r = 0$ contribute.

For any $\epsilon > 0$ by adjusting $\delta, \delta_2, \delta_3$ we can get a state $|\Phi\rangle$ which approximates $|\psi\rangle^{\otimes n}$ with fidelity $1 - \epsilon$.

To proceed we first consider a particular superselection selector specified by a choice of $\{j_1, j_2, \dots, j_n\}$. The component of $|\Phi\rangle$ in such a sector lies in a space of dimension

$$d = \sum_{s=\bar{s}(1-\delta_2)}^{s=\bar{s}(1+\delta_2)} \sum_{t=\bar{t}(1-\delta_3)}^{t=\bar{t}(1+\delta_3)} \binom{m}{s} \binom{n-m}{t}, \quad (4.56)$$

where m is given by eq. (4.55). Importantly, d is the same for all superselection sectors with the same value of m .

As a result, one can consider the set of all superselection sectors which correspond to the same value of m together. Alice can use two sets of q_m qubits which are initially unentangled and in the state

$$|000 \dots 00\rangle \otimes |000 \dots 00\rangle. \quad (4.57)$$

and extract the entanglement in all these superselection sectors. The number of qubits needed are

$$q_m = S_q(m)(1 + O(1/\sqrt{n})) \quad (4.58)$$

where

$$S_q(m) = -m[\alpha_{1,1} \log \alpha_{1,1} + \alpha_{0,1} \log \alpha_{0,1}] - (n-m)[\alpha_{1,0} \log \alpha_{1,0} + \alpha_{0,0} \log \alpha_{0,0}] \quad (4.59)$$

This follows from noting that d defined in eq. (4.56) takes the value

$$d = 2^{S_q(m)(1+O(1/\sqrt{n}))}. \quad (4.60)$$

To extract the entanglement in these superselection sectors Alice swaps the state of the first set of q_m qubits with that of A' in each superselection sector, and then the state of the second set of qubits with C also in the same sector. The second set of qubits is then teleported to Bob using Bell pairs, following which Bob then swaps its state with that of B also in the same superselection sector. Finally, Alice swaps the state of the first set of qubits with that of A also in the same superselection sector. The number of Bell pairs used up in the process is q_m .

Actually, the last two swaps need to be defined more precisely as follows. Bob takes the state of the second set of qubits obtained after teleportation and swaps it with states in a d dimensional subspace of B obtained by taking the quantum numbers s, t defined in eq. (4.53), eq. (4.54), now referring to the system B , in the range, $s \in \{\bar{s}(1 - \delta_2), \bar{s}(1 + \delta_2)\}$, $t \in \{\bar{t}(1 - \delta_3), \bar{t}(1 + \delta_3)\}$. Similarly for the last swap transformation that Alice carries out. Also, we note that by adjusting δ_2, δ_3 (while keeping them to be $O(1/\sqrt{n})$) we can ensure that q_m is an integer.

This procedure can be extended for all values of

$$m \in \{\bar{m}(1 - \delta), \bar{m}(1 + \delta)\} \quad (4.61)$$

in the sum, eq. (4.52), as follows. The number of values m takes is

$$\Delta m = 2^{\Delta q}, \quad (4.62)$$

where $\Delta q \sim O(\log \sqrt{n})$. This follows from eq. (4.49) and eq. (4.50). One can therefore keep track of which value of m one is working with by adding an additional Δq number of pairs of qubits (this number can also be made an integer by adjusting δ).

For this purpose we start with

$$q_T = \Delta q + q_{\max} \quad (4.63)$$

number of pairs of qubits, where q_{\max} is the maximum value $q(m)$ takes as m ranges over the set in eq. (4.61). It follows from eq. (4.49), eq. (4.58) that

$$q_{\max} = nS_q(1 + O(1/\sqrt{n})). \quad (4.64)$$

Each of the Δq pairs can be taken in a Bell pair state. As a result the state of all of these pairs is a symmetric superposition of Δm states, with one set of Δq qubits being

completely correlated with the other set. The remaining q_{\max} pairs are in the states, eq. (4.57). The q_T pairs can then be divided into two sets, with each set containing half of the Δq Bell pairs and one set of the q_{\max} pairs.

Depending on the state of the Δq qubits in the first set Alice then carries out a swap between the state of A' in the set of superselection sectors with the corresponding value of m and the q_{\max} qubits in the first set. And she carries out a swap between C for the same value of m and the second set of q_{\max} qubits. More correctly, Alice carries out a swap between the state of A' with the appropriate value of m and a subset of the state of the q_{\max} qubits of dimension $q(m)$, eq. (4.58), and similarly for C . The second set of q_T qubits is then teleported to Bob using Bell pairs, and then swapped with B . Once again this is done by relating the values of the first Δq qubits to the value of m in the state in possession of Bob, eq. (4.48). And the first set of q_T qubits is then swapped similarly with A .

The total number of Bell pairs consumed in this process is

$$k' = q_T = nS_q + O(\sqrt{n}) + O(\log \sqrt{n}). \quad (4.65)$$

The result of these operations is that $|\psi^{\text{in}}\rangle$, eq. (4.48), turns into the state $|\Phi\rangle$ which is now shared between A and B . From our discussion above it follows that this final state is a good copy of $|\psi\rangle^{\otimes n}$ with fidelity $1 - \epsilon$. By taking $n \rightarrow \infty$ we can take $\epsilon \rightarrow 0$.

The number of Bell pairs consumed in this limit becomes

$$\lim_{n \rightarrow \infty} \frac{k'}{n} = S_q \quad (4.66)$$

which agrees with eq. (4.47) above.

4.4 Concluding comments on the toy model

In the usual discussion of distillation and dilution, without superselection sectors, the end point of distillation is an unentangled state, and so is the starting point of the dilution process. One also finds that the maximum number of Bell pairs produced k during distillation and the minimum number k' used during dilution satisfy the condition

$$k < k' \quad (4.67)$$

with

$$\lim_{n \rightarrow \infty} \frac{k}{n} = \lim_{n \rightarrow \infty} \frac{k'}{n} = S_{EE} \quad (4.68)$$

where S_{EE} is the entanglement entropy. This allows one to conclude that the protocols one has identified are optimal. For if one could have carried out the dilution process while consuming fewer Bell pairs, then combining that with the known protocol for distillation would lead to a process where one began and ended with the same unentangled state but produced some net number of Bell pairs. This would be a contradiction since only LOCC were involved.

In the discussion above with superselection sectors we have seen that k and k' do meet the conditions,⁶ eq. (4.67) and eq. (4.68). However the end point of distillation given in

⁶While we did not elaborate on eq. (4.67) above, it essentially follows from the inequality eq. (4.40) and eq. (4.65).

eq. (4.45) and the state we needed to begin the dilution process, eq. (4.48), are not the same. Thus one might wonder if one has identified the optimal protocols in this case. Perhaps one can improve them and extract more Bell pairs than given in eq. (4.68)?

We now argue that this is not possible and the limit S_q identified above is indeed the best one can do. The point is the following. We had already argued in section 4.1 that using only local gauge-invariant operators we should not be able to extract the full entanglement in the Bipartite state. A residual entanglement given by the S_c term in eq. (4.18) must remain unextracted from the Bipartite state. This expectation is in fact borne out in the two protocols above. It is easy to see that the entanglement per copy in the residual state left over after distillation eq. (4.45) is indeed S_c , as is the entanglement in the starting state, eq. (4.48), of dilution.

Thus, the beginning state of distillation and the end point of dilution in our protocols, while not being the same, do achieve the minimum bound on the residual entanglement that cannot be extracted. It then follows that the maximum entanglement that could have been extracted in Bell pairs is given by S_q , the second term in eq. (4.17), eq. (4.19), which is indeed met by the protocols above.

The argument about S_q being the maximum entanglement that can be extracted can be seen as follows. Suppose we could have extracted more entanglement than S_q . Then, at the end of the distillation process, tasteless as it may seem, working in the extended Hilbert space where we allow non-gauge-invariant operators, we could also extract the entanglement from the residual state which equals S_c . This would result in a total extracted entanglement which is bigger than $S_c + S_q$, eq. (4.17), thereby leading to a contradiction.

4.5 Comments on gauge theories

Let us end by connecting our discussion of the toy model above with that for gauge theories. As was mentioned in section 4.1 above, the S_c term, eq. (4.18), is analogous to the first term in eq. (2.11) in the Abelian case and the first two terms in eq. (2.30) for the Non-Abelian case. We have seen in the toy model that the entanglement corresponding to the S_c term cannot be extracted during distillation or dilution. This suggests that the corresponding terms also cannot be extracted in the Abelian and Non-Abelian cases respectively. It follows then that the maximum entanglement one can extract, in the asymptotic sense where we work with n copies and take $n \rightarrow \infty$, is given by the second term in eq. (2.11),

$$S_{q,A} = - \sum_{\mathbf{k}} p_{\mathbf{k}} \text{Tr}_{\mathcal{H}_{\text{in}}^{\mathbf{k}}} \tilde{\rho}_{\text{in}}^{\mathbf{k}} \log \tilde{\rho}_{\text{in}}^{\mathbf{k}}, \quad (4.69)$$

for the Abelian case and by the third term in eq. (2.30),

$$S_{q,NA} = - \sum_{\mathbf{k}} p_{\mathbf{k}} \text{Tr}_{\hat{\mathcal{H}}_{\text{in}}^{\mathbf{k}}} \tilde{\rho}_{\text{in}}^{\mathbf{k}} \log \left(\tilde{\rho}_{\text{in}}^{\mathbf{k}} \right), \quad (4.70)$$

for the Non-Abelian case. This conclusion is in fact true and follows quite straightforwardly from our discussion of the toy model as we will now see. Eq. (4.69) and eq. (4.70) are some of the main conclusions of this paper. For a more complete definition of the various symbols

appearing in these equations we refer the reader to section 2 especially the discussion around eq. (2.11) and eq. (2.30).

The toy model can in fact be immediately generalised to apply to an Abelian gauge theory. As discussed in section 3, the Hilbert space of gauge-invariant states $\mathcal{H}_{\text{ginv}}$ in the Abelian case can be expressed as a sum over tensor products, eq. (3.6), where the sum is over different superselection sectors labelled by \mathbf{k} .

In the toy model the second and third qubits, in the wave function, eq. (4.4) took the same values and could not be changed by allowed local operators. These qubits then encode for the superselection sectors in the model. To extend the toy model for the Abelian case we simply add more qubits in both A and B , which are enough in number to encode for the full superselection vector \mathbf{k} . The constraint of gauge-invariance is that these qubits must take the same values in A and B . Also we increase the dimension of the Hilbert spaces spanned by the first and fourth qubits in eq. (4.4) to equal the dimension of $\mathcal{H}_{\text{ginv},\text{in}}^{\mathbf{k}}, \mathcal{H}_{\text{ginv},\text{out}}^{\mathbf{k}}$ respectively. Gauge-invariant operators acting on inside links allow states in $\mathcal{H}_{\text{ginv},\text{in}}^{\mathbf{k}}$ to be transformed from one to another in general, and similarly for $\mathcal{H}_{\text{ginv},\text{out}}^{\mathbf{k}}$. Our discussion of the toy model then applies in a straightforward way to the Abelian case and shows that the maximum entanglement which can be extracted is given by eq. (4.69). The residual entanglement left over after distillation, or which must be present at the start of dilution, is given by the first term in eq. (2.11).

The extension to the Non-Abelian case is also straightforward. Recall once again, that in the context of the electric centre definition in section 3, we found that the space of gauge invariant states could be written as a sum over tensor products, with the sum being over different superselection sectors, eq. (3.15), for the Non-Abelian case as well. The only difference here is that the labels of each superselection sector specify the irreducible representation of the Non-Abelian gauge group under which the total ingoing (or outgoing) flux at each boundary vertex transforms. The extension required for the toy model to describe this case is then also very similar to the Abelian case of the previous paragraph. The second and third qubit in the toy model are extended to describe the superselection sectors in the Non-Abelian case, and the dimension of the spaces spanned by the first and fourth qubits are extended to span the spaces $\mathcal{H}_{\text{ginv},\text{in}}^{\mathbf{k}}, \mathcal{H}_{\text{ginv},\text{out}}^{\mathbf{k}}$, which were defined in section 3.2 for the Non-Abelian case. This leads to the conclusion that the maximum entanglement which can be extracted in the Non-Abelian case is given by the second term in eq. (3.21), which is the third term in eq. (2.30) and in agreement with eq. (4.70).

The residual entanglement which cannot be extracted depends on the definition which is used. For the electric centre definition this is given by the first term in eq. (3.21), which is also the first term in eq. (2.30). For the extended Hilbert space definition the residual entanglement is given by the first two terms in eq. (2.30) as was mentioned above.

The extension of the toy model for the Non-Abelian case as described above actually gives the description of the state in $\mathcal{H}_{\text{ginv}}$ which was discussed in section 3.2. The model can be further extended to directly give the description of a state in the extended Hilbert space, \mathcal{H} . Since $\mathcal{H}_{\text{ginv},\text{in}}^{\mathbf{k}}, \mathcal{H}_{\text{ginv},\text{out}}^{\mathbf{k}}$, are already identified with $\hat{\mathcal{H}}_{\text{ginv},\text{in}}^{\mathbf{k}}, \hat{\mathcal{H}}_{\text{ginv},\text{out}}^{\mathbf{k}}$, respectively, eq. (3.13), eq. (3.14), one needs to augment the Hilbert spaces spanned by the second and third qubits of the toy model further by also adding the spaces $\mathcal{H}^1 \otimes \cdots \otimes \mathcal{H}^n$,

and $\mathcal{H}'^1 \otimes \cdots \otimes \mathcal{H}'^n$ respectively to them for this purpose. As was mentioned in the discussion around eq. (3.12), given a state in $\mathcal{H}_{\text{ginv},\text{in}}^{\mathbf{k}} \otimes \mathcal{H}_{\text{ginv},\text{out}}^{\mathbf{k}}$ we can construct a unique state in $\mathcal{H}^{\mathbf{k}}$ from it. Summing over the different superselection sectors \mathbf{k} then gives the state in \mathcal{H} . In this way, after the additional augmentation the toy model directly describes states in the extended Hilbert space. It is then straightforward to see that the analysis above when carried out in this augmented case leads to the conclusion that the entropy which can be extracted is given by eq. (4.70) and the residual entanglement entropy which cannot be extracted is the first two terms in eq. (2.30).

5 Degeneracy of the non-Abelian toric code

Since, we as saw above, the full entanglement entropy given by the extended Hilbert space definition cannot be extracted in distillation or dilution, one might ask whether an alternate definition given by the entanglement which can be extracted in these processes, by the most optimal protocols, is in fact more appropriate.

Here, we will examine a class of Non-Abelian toric codes [14] and find that the extended Hilbert space definition has the attractive feature of leading to the correct topological entanglement entropy, [12, 13]. It will turn out in these models that the extra terms which arise in the extended Hilbert space definition (the first two terms in eq. (2.30)) make an important contribution to the full answer. An alternate definition, corresponding to the number of Bell pairs which can be extracted, will therefore not lead to the correct topological entropy.

In fact, this is already true in the Abelian case, where the full entanglement, eq. (2.11) is needed and the second term alone is not good enough, [2]. Here we will discuss the Non-Abelian case to also bring out the significance of the second term in eq. (2.30) which arises only in this context.

The fact that the extended Hilbert space definition gives the correct answer to the topological entanglement entropy for gauge theories based on toric codes is actually more or less guaranteed to be true from their very construction. In these models gauge-invariance is only imposed “weakly” to begin with, by a term in the Hamiltonian which makes it energetically unfavourable to break this symmetry. While the resulting ground state is gauge-invariant the excited states are therefore not, in general. And the correct entanglement entropy is obviously given by working in the extended Hilbert space. The gauge theory is recovered in the limit in which the electric coupling $J_e \rightarrow \infty$ (this coupling is defined in eq. (5.4) below). It then follows that the natural definition of entanglement in the gauge theory should also be obtained by the extended Hilbert space definition in this limit. Moreover, since the topological entanglement entropy is by construction invariant under smooth deformations of the Hamiltonian, it should not change as this limit is taken (provided correlation lengths continue to be finite, as will be the case here). Thus, the extended Hilbert space definition should lead to the correct topological entanglement entropy in the gauge theory too.

The purpose of carrying out the explicit calculations here is to illustrate that the extra terms present in the extended Hilbert space definition, do contribute in an essential way in the answer for the topological entanglement entropy.

We work in this section with toric codes based on a discrete non-Abelian symmetry group G . The dimension of this group will be denoted by $|G|$, and its irreducible representations by $r_i, i = 1, \dots, n$. A concrete example to keep in mind is S_3 , with dimension 6 and three irreducible representations.

We work in 2 spatial dimensions. The degrees of freedom live on links L_{ij} . For each link these degrees lie in a Hilbert space \mathcal{H}_{ij} of dimension $|G|$. Suppressing the link indices for the moment, a basis of the Hilbert space at each link is given by the orthonormal vectors, $\{|g\rangle : g \in G\}$. The corresponding basis elements for \mathcal{H}_{ij} are denoted by $\{|g\rangle_{ij}\}$. A basis for the full Hilbert space $\mathcal{H} = \otimes_{ij} \mathcal{H}_{ij}$ is obtained by taking the tensor product $\otimes |g\rangle_{ij}$ over all the links. Note that each link L_{ij} has an orientation. If this link is in the state $|g\rangle_{ij}$ then its oppositely oriented counterpart L_{ji} , would be in the state $|g^{-1}\rangle_{ji}$.

There is a left and right action of the group which can be defined by operators $\mathcal{L}_{+,ij}^h, \mathcal{L}_{-,ij}^h$ on each link as follows:

$$\mathcal{L}_{+,ij}^h |g\rangle_{ij} = |h \cdot g\rangle_{ij} \quad (5.1)$$

$$\mathcal{L}_{-,ij}^h |g\rangle_{ij} = |g \cdot h^{-1}\rangle_{ij} \quad (5.2)$$

for any $h \in G$. This is analogous to the action of the operators, $\hat{J}^a, \hat{\mathcal{J}}^a$ in the $SU(2)$ case, eq. (2.12), eq. (2.13).

We also define projectors, $T_{+,ij}^h, T_{-,ij}^h$, by

$$T_{+,ij}^h |g\rangle = \delta_{h,g} |g\rangle_{ij}, \quad T_{-,ij}^h |g\rangle = \delta_{g^{-1},h} |g\rangle_{ij} \quad (5.3)$$

for any $h \in G$.

The Hamiltonian is given by the sum of two terms:

$$H = -J_e \sum_i A_i - J_m \sum_p B(p) \quad (5.4)$$

with i denoting all vertices, V_i , in the lattice and p denoting the sum over all elementary plaquettes. The couplings, $J_m, J_e > 0$.

The operator A_i is defined at each vertex V_i as follows. We take all links L_{ij} to be oriented outward from the vertex, and take,

$$A_i = \sum_g \prod_j \mathcal{L}_{+,ij}^g \quad (5.5)$$

where $\mathcal{L}_{+,ij}^g$, defined in eq. (5.1), acts on the state of the L_{ij} link. The sum in eq. (5.5) is over all elements of G . The product in eq. (5.5) takes values over all links emanating from V_i . Under a gauge transformation, parametrised by group element h , at vertex V_i , the state $|g\rangle_{ij}$ of the Link L_{ij} which emanates from i and goes to j transforms as $|g\rangle_{ij} \rightarrow \mathcal{L}_{+,ij}^h |g\rangle_{ij}$. Thus we see that the sum in A_i corresponds to a sum over all gauge transformations at vertex V_i and the first term in H is then a sum over all gauge transformations for all vertices.

The operator $B(p)$ is defined as follows. Consider the elementary plaquette p and orient the links in the plaquette to run counter clockwise. There are four such links which we label as 1, 2, 3, 4. Then

$$B(p) = \sum_{h_1, h_2, h_3, h_4} T_{+,1}^{h_1} T_{+,2}^{h_2} T_{+,3}^{h_3} T_{+,4}^{h_4}, \quad (5.6)$$




Figure 2. The topological entanglement entropy can be calculated from three regions of this sort, by the formula, eq. (5.9).

where the sum is over all values of group elements h_i, h_2, h_3, h_4 , subject to the condition $h_1 h_2 h_3 h_4 = 1$. Loosely speaking the operator $B(p)$ projects onto states where the magnetic flux around the plaquette is identity.

It is easy to see that the two terms in H commute with each other. This makes it easy to find the ground state, $|s\rangle$. This state is unique on the plane, \mathbb{R}^2 . It is gauge-invariant and satisfies the conditions,

$$A_i |s\rangle = |s\rangle, \quad (5.7)$$

at each vertex, and

$$B(p) |s\rangle = |s\rangle \quad (5.8)$$

for each plaquette. The ground state is given by the symmetric sum of all states for which the magnetic flux around each plaquette takes the value identity. We will be interested in the topological entanglement entropy of $|s\rangle$.

Consider three regions A, B, C shown in figure 2. Let S_A be the entanglement entropy of A with the rest of the system, $S_{A \cup B}$ the entanglement of $A \cup B$ with the rest, etc. Then the topological entanglement is defined as

$$S_{\text{top}} = S_A + S_B + S_C - S_{A \cup B} - S_{B \cup C} - S_{C \cup A} + S_{A \cup B \cup C}. \quad (5.9)$$

To compute it we will first consider a single region which is the set of links on and inside a rectangle as shown in figure 1 and show how its entanglement is computed. The result for S_{top} will then follow.

For the rectangular region let there be n boundary links taking values, $1, 2, \dots, n$, ordered in an anticlockwise fashion. And let us consider a configuration where the degree of freedom on each of these boundary links take values, $\{h_1, h_2, \dots, h_n\}$, meeting the condition,

$$h_1 h_2 \dots h_n = 1 \quad (5.10)$$

as required for state $|s\rangle$. Due to the constraint, eq. (5.10), only $n - 1$ of the link variables are independent resulting in $|G|^{n-1}$ such independent configurations. Now since the magnetic flux on all plaquettes is completely determined in the state $|s\rangle$, eq. (5.8), it can be

shown that the resulting state which contributes to the ground state $|s\rangle$ with the choice $\{h_1, h_2, \dots, h_n\}$ has the form,

$$|h_1 h_2 h_3 \dots h_n\rangle \otimes |\psi_{\{h_i\}, \text{out}}\rangle \otimes |\psi_{\{h_i\}, \text{in}}\rangle. \quad (5.11)$$

Here $|\psi_{\{h_i\}, \text{out}}\rangle, |\psi_{\{h_i\}, \text{in}}\rangle$, are pure states defined on the Hilbert spaces of the outside and inside links, and $|h_1 h_2 h_3 \dots h_n\rangle$ is a shorthand for the state $|h_1\rangle \otimes |h_2\rangle \otimes \dots \otimes |h_n\rangle$ of the boundary links. Further, the states $|\psi_{\{h_i\}, \text{out}}\rangle$, in eq. (5.11), for distinct choices of the boundary links, $\{h_1, h_2, \dots, h_n\}$, are orthogonal. Similarly, the states $|\psi_{\{h_i\}, \text{in}}\rangle$ are also orthogonal.⁷

The state $|s\rangle$ can then be written down as the sum of all such terms,

$$|s\rangle = \frac{1}{|G|^{\frac{n-1}{2}}} \sum_{\{h_i\}} |h_1 h_2 h_3 \dots h_n\rangle \otimes |\psi_{\{h_i\}, \text{out}}\rangle \otimes |\psi_{\{h_i\}, \text{in}}\rangle, \quad (5.12)$$

where the sum is over all values of $\{h_1, h_2, \dots, h_n\}$ subject to the condition eq. (5.10). There are $|G|^{n-1}$ orthogonal states in the sum resulting in the pre factor in eq. (5.12) which makes $|s\rangle$ of unit norm.

Tracing over the outside links then gives the density matrix

$$\rho_{\text{in}} = \frac{1}{|G|^{n-1}} \sum_{\{h_i\}} |h_1 h_2 h_3 \dots h_n\rangle \otimes |\psi_{\{h_i\}, \text{in}}\rangle \langle h_1 h_2 h_3 \dots h_n| \otimes \langle \psi_{\{h_i\}, \text{in}}| \quad (5.13)$$

Since the density matrix is diagonal in the basis, $|h_1 h_2 \dots h_n\rangle \otimes |\psi_{\{h_i\}, \text{in}}\rangle$, in fact a multiple of the identity, the entanglement is easily computed. We get,

$$S_{EE} = (n-1) \log(|G|) \quad (5.14)$$

due to the $n-1$ independent boundary links.

The topological entanglement can now be easily calculated for the three regions shown in figure 2 and gives

$$S_{\text{top}} = -\log(|G|) \quad (5.15)$$

5.1 The different superselection sectors

Let us now go back to the rectangular region in figure 1 and try to understand our result for the entanglement in terms of contributions from different superselection sectors as defined in section 2. The links of interest are those inside and on the boundary of the rectangle, their degrees of freedom lie in \mathcal{H}_{in} . We will refer to the basis $\{|h_1 h_2 \dots h_n\rangle\}$ used to describe the states of the boundary links above as the “magnetic basis” below. The states $|h\rangle$ of a link are not gauge-invariant under a gauge transformation acting on its end points. This means that the basis $\{|h_1 h_2 \dots h_n\rangle\}$ is not invariant under boundary gauge transformations. By a boundary gauge transformation we mean a gauge transformation acting at one of the boundary vertices.

The basis in which the contribution from different superselection sectors becomes manifest is a different one. It is obtained by decomposing the Hilbert space of the inside links,

⁷Eq. (5.11) can be seen, for example, by fixing gauge at all vertices, other than boundary vertices. Such a gauge fixing does not change the entanglement entropy.

\mathcal{H}_{in} , into irreducible representations of the boundary gauge transformations. Although the basis elements $|h_1 h_2 \cdots h_n\rangle$ for the boundary links are not invariant under these transformations, an important point to note is that the sector satisfying the condition, eq. (5.10), for which the density matrix has non-trivial support, is indeed invariant. This sector can therefore by itself be decomposed into irreducible representations of the boundary gauge transformations. Notice also, as is clear from eq. (5.12), eq. (5.13), that in this sector the density matrix has a very simple form, namely being a multiple of the identity. It therefore must continue to have this form even when we decompose the sector into the irreducible representations of the boundary gauge transformations.

Our job of expressing the entanglement in terms of different superselection sectors is therefore quite easy. We have simply to figure out which irreducible representations of the boundary gauge transformations occur after we carry out this basis change.

A superselection sector is specified by the vector $\mathbf{k} = \{r_1, r_2, \cdots, r_n\}$. States of \mathcal{H}_{in} lying in this sector transform as the r_i representation under a boundary gauge transformation at the vertex V_i . To see why this specifies a superselection sector, we note that Gauss' law then tells us that the outside links (oriented suitably) leaving V_i must transform as the conjugate representation, \bar{r}_i . Now, the state of the outside links cannot be changed by operators acting on the inside links alone. Thus the superselection sectors cannot be changed by gauge-invariant operators acting only on the inside region.

It is easy to see that the different representations r_1, r_2, \cdots, r_n which enter in \mathbf{k} are not all independent. The constraint eq. (5.10) satisfied by the boundary links leads to a condition amongst them. For given values $r_1, r_2, \cdots, r_{n-1}$ of the representations under the gauge transformations at the first $n - 1$ boundary vertices one finds that the only values for r_n which appear are those which can combine with the others to give a singlet of G . For the state $|s\rangle$ we started with, this is in fact the only constraint. Since the state is a symmetric sum over all links satisfying the condition eq. (5.10), all values of $\{r_1, r_2, \cdots, r_n\}$ must appear, subject to this one constraint, when we decompose the sector, eq. (5.10), into irreducible representations. While we omit some of the steps here, this result can be obtained by starting with the basis $|h_1, h_2, \cdots, h_n\rangle$ for the boundary links and explicitly changing to a new basis which transforms as an irreducible representation under the boundary gauge transformations.

As a check we note the following. Let us denote the dimension of the r_i^{th} representation of G to be d_i , so that the r_1 representation has dimension d_1 , etc. Then the dimension of the representation obtained by taking a tensor product of the, $\{r_1, r_2, \cdots, r_n\}$ representations is $d_1 d_2 \cdots d_n$. For a given set of values of the first $n - 1$ labels, $\{r_1, r_2, \cdots, r_{n-1}\}$ we get a subspace of dimension

$$d_1 d_2 \cdots d_{n-1}. \quad (5.16)$$

For the state to be a singlet under global gauge transformations, r_n must take the values corresponding to all the irreducible representations that appear in $r_1 \otimes \cdots \otimes r_{n-1}$. Therefore, for any given set of values $\{r_1, \cdots, r_{n-1}\}$ the set of values r_n can take must satisfy

$$\sum_{r_n} d_n = d_1 \cdots d_{n-1}. \quad (5.17)$$

However, a representation may appear in multiple linearly independent ways in the tensor product of $n-1$ representations; this corresponds to the fact that the identity representation appears in multiple linearly independent ways in the tensor product $r_1 \otimes \cdots \otimes r_n$. Since the superselection sectors are completely specified by the representations $\{r_1, \cdots r_n\}$, all the combinations belong to the same superselection sectors.⁸ We will denote the number of combinations as $N_{\mathbf{k}}$ below. In terms of our discussion in section 2.2, the subspace of $\mathcal{H}_{\text{in}}^{\mathbf{k}}$, the subspace of \mathcal{H}_{in} corresponding to the superselection sector \mathbf{k} , on which $\rho_{\text{in}}^{\mathbf{k}}$ has support is of the form given in eq. (2.22), with $\mathcal{H}^i, i = 1, \cdots n$, having dimension d_i , and $\hat{\mathcal{H}}_{\text{in}}^{\mathbf{k}}$ being replaced by a subspace of dimension $N_{\mathbf{k}}$.

Summing over all values of $r_1, \cdots r_n$, with this one constraint then gives the total dimension, to be

$$N_T = \sum_{r_1, r_2, \cdots r_n} d_1 d_2 \cdots d_n = \sum_{r_1, r_2, \cdots r_{n-1}} d_1^2 d_2^2 \cdots d_{n-1}^2 = |G|^{n-1}, \quad (5.18)$$

where we have used eq. (5.17), and the relation

$$\sum_r d_r^2 = |G|. \quad (5.19)$$

We see that N_T is in fact equal to the total dimension of the space of states which occur in $|s\rangle$, eq. (5.12).

It was mentioned above that after the change of basis ρ_{in} must continue to be a multiple of the identity. From this it follows that the probability to be in any superselection sector, $p_{\mathbf{k}}$, is proportional to the number of the states in the superselection sector for which ρ_{in} has non-trivial support, resulting in,

$$p_{\mathbf{k}} = \frac{(\prod_{i=1}^n d_i) N_{\mathbf{k}}}{|G|^{n-1}}. \quad (5.20)$$

The entanglement entropy is then given by

$$S_{EE} = - \sum_{\mathbf{k}} p_{\mathbf{k}} \log p_{\mathbf{k}} + \sum_{\mathbf{k}} p_{\mathbf{k}} \left(\sum_{i=1}^n \log d_i \right) + \sum_{\mathbf{k}} p_{\mathbf{k}} \log N_{\mathbf{k}}. \quad (5.21)$$

The three terms on the r.h.s. correspond to the three terms in eq. (2.30) respectively. In particular, while first two terms cannot be extracted in distillation or dilution, the last term can be measured through these processes. It is easy to see from eq. (5.18), eq. (5.19) that the entanglement in eq. (5.21) agrees with eq. (5.14).

Now that we have verified that the full answer for the entanglement in the ground state of the toric codes considered here gets a significant contribution from terms which are not amenable to extraction in distillation and dilution, it also follows that dropping these terms will not give rise to the correct topological entanglement. Let us also note that the topological entanglement entropy on general grounds is related to the total quantum

⁸The original version of the paper had an error. It claimed that only one singlet state could appear in each superselection sector. The discussion, with the correct argument given here appeared in [22].

dimension, [12, 13]. From this general relation and our result, eq. (5.15) we see that total quantum dimension, \mathcal{D} , in the models we have been considering here is given by

$$\mathcal{D} = |G| \quad (5.22)$$

It is easy to verify that this is the correct answer in simple examples. E.g. for S_3 , we get $\mathcal{D} = 6$, which agrees with the value obtained after summing over the quantum dimensions of all particles in the model (see, e.g., [23]).

6 Entanglement entropy in the strong-coupling expansion

We have seen in the section above on the Non-Abelian toric code that the entanglement in the ground state arises due to the contribution from the first two terms in eq. (2.30). Here, we study an $SU(2)$ gauge theory in the strong-coupling expansion. Working to first non-trivial order we find once again that the entanglement arises entirely due to the first two terms in eq. (2.30). This calculation has been done earlier [4, 15, 24], but we reproduce it here to emphasise the role of these terms.⁹

The set up here is closely related to that in section 2.2 where we also discussed the $SU(2)$ theory, and we use the same notation. We work on a square lattice. The degrees of freedom on each link L_{ij} lie in a Hilbert space \mathcal{H}_{ij} with states, $|U_{ij}\rangle$. The Hamiltonian of this model, see [26], after suitably rescaling, is

$$H = \frac{g^2}{2} \sum_{(ij)} \hat{J}_{ij}^2 + \sum_{(ijkl)} \frac{4}{g^2} \text{tr} \left(\hat{P}_{\{ijkl\}} + \hat{P}_{\{ijkl\}}^\dagger \right), \quad (6.1)$$

where

$$\hat{P}_{\{ijkl\}} = \hat{U}_{ij} \hat{U}_{jk} \hat{U}_{kl} \hat{U}_{li} \quad (6.2)$$

is the product of all link variables along an elementary plaquette formed by the links, $L_{ij}, L_{jk}, L_{kl}, L_{li}$. The sum in the first term is over all links (ij) , and in the second term over all elementary plaquettes $(ijkl)$. The operator $\hat{J}_{ij}^2 = \sum_{a=1}^3 (\hat{J}_{ij}^a)^2$ was described in section 2.2, eq. (2.12). Each link appears once in the sum in the first term in eq. (6.1). Since $\hat{J}_{ij}^2 = \hat{J}_{ji}^2$, eq. (2.14), eq. (2.15), the orientation of the link does not matter.

g^2 in eq. (6.1) is the coupling constant. We work in the strong coupling limit where $g^2 \gg 1$. The Hamiltonian eq. (6.1) can also be written as

$$H = \frac{g^2}{2} \left[H^0 + \frac{3}{2} \epsilon H^1 \right], \quad (6.3)$$

where

$$H^0 = \sum_{(ij)} \hat{J}_{ij}^2, \quad (6.4)$$

$$H^1 = \text{tr} \left(\hat{P}_{\{ijkl\}} + \hat{P}_{\{ijkl\}}^\dagger \right), \quad (6.5)$$

⁹For an interesting discussion of the relation of this answer and that of the previous section to confinement, see [15, 25].

$$\epsilon = \frac{16}{3g^4}. \quad (6.6)$$

The prefactor $\frac{g^2}{2}$ can be absorbed in H by a rescaling giving

$$H = H^0 + \frac{3}{2}\epsilon H^1. \quad (6.7)$$

In the strong coupling limit, $\epsilon \ll 1$. We will be interested in computing the ground state of H in perturbation theory in ϵ and then computing the entanglement of a rectangular region, shown in figure 1. We include the links on the boundary in our choice of the inside set of links.

To leading order the Hamiltonian is given by H^0 and the ground state is an eigenstate with zero angular momentum for all links, i.e., $(J_{ij})^2 = 0$ for all links L_{ij} . We denote the link L_{ij} in the zero angular momentum state by $|0\rangle_{ij}$ and the full ground state, which is a tensor product, $\otimes_{(ij)} |0\rangle_{ij}$, by $|0^{(0)}\rangle$.

To construct the first order correction to the ground state wave function, we note that the operator $\text{tr}(\hat{P}_{\{ijkl\}})$ acting on $|0^{(0)}\rangle$ excites the links appearing in the plaquette to a state which now carries 1/2 units of angular momentum. This follows from the commutation relations,

$$[\hat{J}_{ij}^a, \hat{U}_{\alpha\beta}] = \frac{-\sigma_{\alpha\gamma}^a}{2} \hat{U}_{\gamma\beta}, \quad (6.8)$$

$$[\hat{\mathcal{J}}_{ij}^a, \hat{U}_{\alpha\beta}] = \hat{U}_{\alpha\gamma} \frac{\sigma_{\gamma\alpha}^a}{2}, \quad (6.9)$$

where $\sigma^a, a = 1, 2, 3$ are the usual Pauli matrices. We also note that $\hat{\mathcal{J}}_{ij}^a = \hat{J}_{ji}^a$, eq. (2.14). Each link L_{ij} has two ends and we can think of \hat{J}_{ij}^a and \hat{J}_{ji}^a as generating an SU(2) transformation at the i and j vertices respectively. From the commutation relations above we see that the state $\hat{U}_{ij}|0\rangle_{ij}$ transforms as a spin 1/2 representation under both of these transformations. The trace in H^1 , eq. (6.5), gives rise to a contraction of indices, and as a result the excitations of two links ending at a vertex of the plaquette combine to give net zero angular momentum with respect to the SU(2) transformations at the vertex. This results in a unique state for the four links, which we denote as $|P_{\{ijkl\}}\rangle$ leading to

$$\text{tr}(\hat{P}_{\{ijkl\}})|0\rangle_{\{ijkl\}} = |P_{\{ijkl\}}\rangle, \quad (6.10)$$

where $|0\rangle_{ijkl}$ stands for the tensor product $|0\rangle_{ij} \otimes |0\rangle_{jk} \otimes |0\rangle_{kl} \otimes |0\rangle_{li}$ of the 4 links each in the state with zero angular momentum.

Acting with $\text{tr}(\hat{P}_{\{ijkl\}}^\dagger)$ we get the same state,

$$\text{tr}(\hat{P}_{\{ijkl\}}^\dagger)|0\rangle_{ijkl} = |P_{ijkl}\rangle. \quad (6.11)$$

It follows from standard time independent perturbation theory then that the first order state is given by

$$|0^{(1)}\rangle = \frac{3}{2}\epsilon \mathcal{P} \frac{1}{E^0 - H^0} \mathcal{P} H^{(1)} |0^{(0)}\rangle, \quad (6.12)$$

where \mathcal{P} is the projector onto the subspace of non-zero energies of H^0 . The full state to this order is then

$$|0\rangle = \mathcal{N} \left[|0^{(0)}\rangle + \frac{3}{2}\epsilon \mathcal{P} \frac{1}{E^0 - H^0} \mathcal{P} H^{(1)} |0^{(0)}\rangle \right], \quad (6.13)$$

where \mathcal{N} is a normalisation to ensure that $|0\rangle$ has unit norm.

It is easy to see that all states excited by a single plaquette insertion of the form eq. (6.10), eq. (6.11), have an eigenvalue, 3, with respect to H^0 , while $E^0 = 0$. This gives,

$$|0\rangle = \mathcal{N} \left[|0^{(0)}\rangle - \frac{\epsilon}{2} H^{(1)} |0^{(0)}\rangle \right]. \quad (6.14)$$

Now there are three kinds of plaquettes, those which have links lying completely outside the region of interest, those which have links lying completely inside, and those which have some inside and outside links (note that as per our definition of the region of interest, the links on the boundary of the rectangle in figure 1 are also in the inside set). We will refer to these as the outside, inside and boundary plaquettes respectively below.

This gives,

$$|0\rangle = \mathcal{N} \left[|0^{(0)}\rangle - \epsilon \left(\sum_{\text{out}} |P_{\{ijkl\}}\rangle_{\text{out}} + \sum_{\text{in}} |P_{\{ijkl\}}\rangle_{\text{in}} + \sum_{\text{bd}} |P_{\{ijkl\}}\rangle_{\text{bd}} \right) \right] \quad (6.15)$$

where the suffixes out, in and bd stand for states obtained by acting with the outside, inside and boundary plaquettes respectively.

Let $n_{\text{in}}, n_{\text{out}}, n_{\text{bd}}$ be the number of in, out and boundary plaquettes respectively, so that

$$N = n_{\text{in}} + n_{\text{out}} + n_{\text{bd}} \quad (6.16)$$

is the total number of plaquettes. Also, a simple calculation shows that the norm of each state $|P_{\{ijkl\}}\rangle$ is

$$\langle P_{\{ijkl\}} | P_{\{ijkl\}} \rangle = 1. \quad (6.17)$$

Then we have that the normalisation is

$$\mathcal{N} = \frac{1}{\sqrt{1 + N\epsilon^2}}. \quad (6.18)$$

It is then easy to see that the density matrix for the inside links that we get, in the extended Hilbert space definition, is

$$\begin{aligned} \rho_{\text{in}} = & \frac{1}{1 + N\epsilon^2} \left[\left\{ |0\rangle_{\text{in}} - \epsilon\sqrt{n_{\text{in}}} \left(\frac{1}{\sqrt{n_{\text{in}}}} \sum_{\text{in}} |P_{\{ijkl\}}\rangle \right) \right\} \left\{ \langle 0|_{\text{in}} - \epsilon\sqrt{n_{\text{in}}} \left(\frac{1}{\sqrt{n_{\text{in}}}} \sum_{\text{in}} \langle P_{\{ijkl\}}| \right) \right\} \right. \\ & \left. + n_{\text{out}}\epsilon^2 |0\rangle_{\text{in}} \langle 0|_{\text{in}} + \epsilon^2 \sum_{\text{bd}} \text{Tr}_{\mathcal{H}_{\text{out}}} |P_{\{ijkl\}}\rangle \langle P_{\{ijkl\}}| \right] \end{aligned} \quad (6.19)$$

$|0\rangle_{\text{in}}$ stands for the state of the inside links obtained by taking the tensor product of the $|0\rangle_{ij}$ states with vanishing angular momentum at each link, and the symbols, $\sum_{\text{in}}, \sum_{\text{bd}}$, stand for a sum over all inside and boundary plaquettes. The first term above arises after tracing over a state in which the outside links are unexcited and have vanishing angular

momentum. The second term after tracing out states in which outside links lying in one outside plaquette have been excited, while the third term arises from states in which a boundary plaquette has been excited.

In terms of different superselection sectors, we see that the contribution in the first line in eq. (6.19) and the first term in the second line correspond to the superselection sector where the total angular momentum at each boundary vertex vanishes. The second term on the second line arises from sectors where the angular momentum at two adjacent boundary vertices, connected by the boundary plaquette appearing in the sum, does not vanish. We see from figure 1, that for the rectangular region under consideration each boundary plaquette has three links in the outside and one link in the inside (since boundary links are also in the inside set). The state $|P_{ijkl}\rangle$, involving a boundary plaquette therefore has three links outside and one link on the boundary of the rectangular region excited with $1/2$ unit of angular momentum. The two boundary vertices with non-zero angular momentum are the ones on which the boundary link of this plaquette ends. The angular momentum, J_T^2 , eq. (2.18), at both these vertices is $3/4$ corresponding to $j = 1/2$. For n boundary links we get n different superselection sectors of this type.

The entanglement entropy can be calculated by adding the contributions from each superselection sector. The probability to be in a sector where the electric flux in two adjacent boundary vertices is non-vanishing can be obtained by taking the trace of ρ_{in} in this sector. We get

$$p_{(1/2,1/2)} = \mathcal{N}^2 \epsilon^2 \approx \epsilon^2. \quad (6.20)$$

There are n_{bd} such sectors. We have used the subscript $(1/2, 1/2)$, to remind ourselves that two boundary vertices have $j = 1/2$. The precise boundary vertices are not important, since we get the same probability for all sectors of this kind.

The probability to be in the sector where the angular momentum at each boundary vertex vanishes is given by taking the trace of the first two terms and given by

$$p_0 = \mathcal{N}^2 [1 + \epsilon^2 (n_{\text{in}} + n_{\text{out}})] \approx 1 - \epsilon^2 n_{\text{bd}}. \quad (6.21)$$

As a check we see that $n_{\text{bd}} p_{(1/2,1/2)} + p_0 = 1$ as required.

Coming back to the sectors which carry non-trivial total angular momentum it is easy to see that the resulting vector \mathbf{k} , defined in section 2.2 has two non-zero values corresponding to the two adjacent boundary vertices with non-zero angular momentum, and these take values $3/4$, eq. (2.21). The Hilbert space in the sector $\mathcal{H}_{\text{in}}^{\mathbf{k}}$ is of the form eq. (2.22), where $n = n_{\text{bd}}$ and only two of the Hilbert spaces $\mathcal{H}^1, \dots, \mathcal{H}^n$ have dimension 2 while the others have dimension 1. Also, it is easy to see that the state in $\hat{\mathcal{H}}_{\text{in}}^{\mathbf{k}}$ is pure. The density matrix in this sector $\rho_{\text{in}}^{\mathbf{k}}$ is also of the form eq. (2.23), where $\hat{\rho}_{\text{in}}^{\mathbf{k}}$ is a number.

The resulting contribution from these sectors to S_{EE} then comes only from the first two terms in eq. (2.29) and is

$$\delta_{(1/2,1/2)} S_{EE} = n_{\text{bd}} [-p_{(1/2,1/2)} \log p_{(1/2,1/2)} + p_{(1/2,1/2)} \log(4)]. \quad (6.22)$$

For the sector where no angular momentum is excited at the boundary vertices we get a contribution to S_{EE} from the first term in eq. (2.29), and potentially the last term. The

reduced density matrix is

$$\frac{1}{1 + (n_{\text{in}} + n_{\text{out}})\epsilon^2} \begin{pmatrix} 1 + n_{\text{out}}\epsilon^2 & \sqrt{n_{\text{in}}}\epsilon \\ \sqrt{n_{\text{in}}}\epsilon & n_{\text{in}}\epsilon^2 \end{pmatrix} \quad (6.23)$$

in the basis $\{|0\rangle_{\text{in}}, \frac{1}{\sqrt{n_{\text{in}}}} \sum |P_i^{\text{in}}\rangle\}$. This matrix has eigenvalues of order ϵ^4 , and therefore its contribution up to $O(\epsilon^2)$ is negligible. The contribution from this sector is then

$$\delta_{(0)} S_{EE} = -p_0 \log p_0. \quad (6.24)$$

Summing, the total entanglement is

$$S_{EE} = n_{\text{bd}}[-p_{1/2,1/2} \log p_{1/2,1/2} + p_{1/2,1/2} \log(4)] - p_0 \log p_0 \approx n_{\text{bd}}\epsilon^2 \left[\log \frac{1}{\epsilon^2} + 3 \right] \quad (6.25)$$

where we have set $\log 4 = 2$. From eq. (6.20) and eq. (6.21) we see that this is of order $\epsilon^2 \log(\epsilon^2)$.

A few more comments are in order. First, we see that the contribution up to the first non-trivial order comes entirely from the first two terms in eq. (2.29), as was mentioned above. Second, since our leading answer is, up to a log correction, of $O(\epsilon^2)$, the reader may wonder whether we should have calculated the corrected wave function to second order in perturbation theory. The change in normalisation to this order has already been taken into account by the factor \mathcal{N} . It is easy to see that additional corrections are not needed. Such a correction which we denote as $|0^{(2)}\rangle$ would contribute at this order to the density matrix only through terms of the form $|0^{(0)}\rangle\langle 0^{(2)}| + |0^{(2)}\rangle\langle 0^{(0)}|$, where the $|0^{(2)}\rangle$ state is orthogonal to $|0^{(0)}\rangle$. As a result, these terms, which only contribute to the superselection sector with zero angular momentum entering at each boundary vertex, give a contribution of $O(\epsilon^4)$ which can be neglected.

Finally, the lattice theory at strong coupling is well known to be confining. This is related to the fact that leading behaviour in eq. (6.25) is proportional to the number of boundary plaquettes, n_{bd} . Since n_{bd} also equals the number of boundary links, n , the entanglement is proportional to the length of the boundary. It is interesting to compare these results with the toric code model studied in section 5. Confinement, in the strong coupling theory, is accompanied with electric order. Electric flux tubes cost energy to excite. As a result Gauss' law is implemented at the boundary in a local way, with the electric flux which enters from a boundary vertex exiting from an adjacent vertex. This results in the leading behaviour being proportional to the number of boundary links and the topological entanglement entropy of the three regions shown in figure 2 vanishing. In contrast, in the toric code there is a global constraint, that the representations $\{r_1, r_2, \dots, r_n\}$ must be able to combine into a singlets, which is the analogue of a global Gauss law. The constraint is a global one since the cost for electric flux tubes to stretch across the inside is not significant. As a result the electric flux on boundary vertices can take all values subject to one overall constraint. This one global constraint results in the entanglement being proportional to $n - 1$, eq. (5.14), leading to a non-trivial topological entanglement.

7 Conclusions

In this paper we explored the extended Hilbert space definition for the entanglement entropy of gauge theories in some detail. The entropy can be expressed as a sum over different superselection sectors. These sectors correspond to different values for the electric flux, specified in a gauge-invariant way, which can enter the links of interest at each of the boundary vertices. The resulting answer, in each sector, gets a contribution from two kinds of terms. The first kind is a sort of classical contribution which arises from the probability to be in this sector. The second kind is a quantum contribution due to quantum correlations in the sector. In the Non-Abelian case the classical contribution includes an extra term which arises as follows. In each sector with nonzero electric flux at a boundary vertex, the set of inside and outside links terminating at this vertex must transform in a non-trivial representation of the gauge group. Due to gauge-invariance these non-trivial representations for the inside and outside links then combine and appear in the wave function only as a singlet of the gauge symmetry. This leads to additional entanglement between the inside and outside links in the Non-Abelian case since the irreducible representations are of dimension greater than one, leading to the extra term in the classical contribution. The result for the entanglement entropy in the extended Hilbert space definition is given in eq. (2.30). The classical terms are the first two on the r.h.s. in eq. (2.30), of these the second term is the one which only arises in the Non-Abelian case.

We also examined the processes of entanglement distillation and dilution for gauge theories in some detail and showed that the number of Bell pairs which can be extracted in these processes are equal, in an asymptotic sense, with a large number of copies of the system, to only the quantum type of contribution, i.e. the third term in eq. (2.30). We argued on general grounds that the quantum contribution was the maximum bound which one could hope to achieve, since local operations involving gauge-invariant operators cannot change the superselection sectors. And we also exhibited protocols for distillation and dilution where the number of extracted or consumed Bell pairs achieved this bound. The operational definition in terms of the number of Bell pairs one can extract therefore gives a smaller value for the entanglement entropy than the extended Hilbert space definition.

An electric centre definition, following [2] in the Abelian case, was also presented for Non-Abelian theories in the paper. It was found, interestingly, that this too differed from the extended Hilbert space definition. The difference arises only for the Non-Abelian case in the second term in eq. (2.30) which is absent in the electric centre definition. This result is therefore “midway” between the extended Hilbert space definition and the operational measure described in the previous paragraph: it contains the first and third terms in eq. (2.30) but not the second term.

These differences between various definitions seem bewildering. Which definition should one trust?

One reason that the entanglement entropy is useful is that it often provides answers to physical questions. Since the questions are physical the answers are unambiguous. For example, the entanglement entropy can be used to define the topological entanglement entropy which in turn is often related to the degeneracy of ground states on a manifold of

non-trivial topology or the quantum dimension of excitations in the system. We calculated the topological entanglement entropy using the extended Hilbert space definition for a class of Toric code models based on discrete Non-Abelian gauge groups here and showed that it gives the correct result for the quantum dimension. The classical terms make a significant contribution to the full answer in these models, including an important contribution coming from the second term in eq. (2.30) which arises only in the Non-Abelian case. Dropping some or all of these classical contributions would therefore not yield the correct result.

These investigations suggest to us that the extended Hilbert space definition is indeed a useful one to use. That it does not agree with the operational measure of the number of Bell pairs that can be extracted from the system may be viewed as a distinctive feature of gauge theories. This property is closely tied to the fact that non-local excitations corresponding to Wilson loops and electric flux loops are present in these theories. It seems important to us to study this universal feature of gauge theories further.

There are many other ways in which this work can be extended. While we did not comment on gauge theories with matter explicitly in the paper, it is in fact easy to apply the extended Hilbert space definition for such systems as well. In this case the matter degrees of freedom live on vertices of the lattice, and the system of interest, the “in” system, is then specified by a state on a set of links and vertices. The extended Hilbert space consists of the space obtained by taking the tensor product of link Hilbert spaces for the gauge field degrees of freedom and vertex Hilbert spaces for matter degrees of freedom. Like in the pure gauge field case, this Hilbert space too by definition admits a tensor product structure. Thus tracing over the outside degrees of freedom, which are in the complement of the inside degrees, we can compute ρ_{in} , eq. (2.3), and then the entanglement entropy, eq. (2.4).

One of the properties which makes the extended Hilbert space definition attractive is that it agrees with a path integral calculation which implements the replica trick. It would be useful to study the continuum limit of the lattice system we have investigated here and analyse the entanglement entropy in this limit. On the lattice it is easy to see that the extended Hilbert space definition is unchanged if one fixes the gauge for vertices securely in the inside or outside. However, gauge fixing the vertices which lie on the boundary, where some links inside and outside terminate, in general changes the result. Going to the continuum, starting from this lattice definition, should help us clarify what kind of gauge fixing is allowed in the continuum too. One expects the results for finite quantities, related to mutual information or topological entanglement entropy for example, should be more universal in the continuum limit.

Finally, in 2 space dimensions it is well known that gauge theories are dual to spin systems. The duality is non-local in space-time. It would be useful to understand how the entanglement entropy transforms under this duality, both in the lattice and in the continuum.

We leave these and other related fascinating questions for the future.

Acknowledgments

We acknowledge discussion with Sumit Das, Rahul Dandekar, Marina Huerta, Gowtham Raghunath Kurri, Gautam Mandal, Shiraz Minwalla, Pratyush Nath, Pranjal Nayak, Hiroshi

Ooguri, Djordje Radičević, Sarath Sankar, Ashoke Sen, Tadashi Takayanagi and Spenta Wadia. We especially thank Sudip Ghosh for several discussions and initial collaboration. SPT thanks the organisers of the “International Workshop on Condensed Matter Physics and AdS/CFT”, held at IPMU, Japan, from May 25-29, 2015, and acknowledges the J. C. Bose Fellowship, Department of Science and Technology, Government of India. We acknowledge support from Department of Atomic Energy, Government of India. Most of all, we thank the people of India for generously supporting research in string theory.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] S. Ghosh, R.M. Soni and S.P. Trivedi, *On the entanglement entropy for gauge theories*, *JHEP* **09** (2015) 069 [[arXiv:1501.02593](https://arxiv.org/abs/1501.02593)] [[INSPIRE](#)].
- [2] H. Casini, M. Huerta and J.A. Rosabal, *Remarks on entanglement entropy for gauge fields*, *Phys. Rev. D* **89** (2014) 085012 [[arXiv:1312.1183](https://arxiv.org/abs/1312.1183)] [[INSPIRE](#)].
- [3] P.V. Buividovich and M.I. Polikarpov, *Entanglement entropy in gauge theories and the holographic principle for electric strings*, *Phys. Lett. B* **670** (2008) 141 [[arXiv:0806.3376](https://arxiv.org/abs/0806.3376)] [[INSPIRE](#)].
- [4] W. Donnelly, *Decomposition of entanglement entropy in lattice gauge theory*, *Phys. Rev. D* **85** (2012) 085004 [[arXiv:1109.0036](https://arxiv.org/abs/1109.0036)] [[INSPIRE](#)].
- [5] H. Casini and M. Huerta, *Entanglement entropy for a Maxwell field: numerical calculation on a two dimensional lattice*, *Phys. Rev. D* **90** (2014) 105013 [[arXiv:1406.2991](https://arxiv.org/abs/1406.2991)] [[INSPIRE](#)].
- [6] D. Radičević, *Notes on entanglement in Abelian gauge theories*, [arXiv:1404.1391](https://arxiv.org/abs/1404.1391) [[INSPIRE](#)].
- [7] W. Donnelly, *Entanglement entropy and non-Abelian gauge symmetry*, *Class. Quant. Grav.* **31** (2014) 214003 [[arXiv:1406.7304](https://arxiv.org/abs/1406.7304)] [[INSPIRE](#)].
- [8] L.-Y. Hung and Y. Wan, *Revisiting entanglement entropy of lattice gauge theories*, *JHEP* **04** (2015) 122 [[arXiv:1501.04389](https://arxiv.org/abs/1501.04389)] [[INSPIRE](#)].
- [9] S. Aoki, T. Iritani, M. Nozaki, T. Numasawa, N. Shiba and H. Tasaki, *On the definition of entanglement entropy in lattice gauge theories*, *JHEP* **06** (2015) 187 [[arXiv:1502.04267](https://arxiv.org/abs/1502.04267)] [[INSPIRE](#)].
- [10] W. Donnelly and A.C. Wall, *Entanglement entropy of electromagnetic edge modes*, *Phys. Rev. Lett.* **114** (2015) 111603 [[arXiv:1412.1895](https://arxiv.org/abs/1412.1895)] [[INSPIRE](#)].
- [11] W. Donnelly and A.C. Wall, *Geometric entropy and edge modes of the electromagnetic field*, [arXiv:1506.05792](https://arxiv.org/abs/1506.05792) [[INSPIRE](#)].
- [12] A. Kitaev and J. Preskill, *Topological entanglement entropy*, *Phys. Rev. Lett.* **96** (2006) 110404 [[hep-th/0510092](https://arxiv.org/abs/hep-th/0510092)] [[INSPIRE](#)].
- [13] M. Levin and X.-G. Wen, *Detecting topological order in a ground state wavefunction*, *Phys. Rev. Lett.* **96** (2006) 110405 [[cond-mat/0510613](https://arxiv.org/abs/cond-mat/0510613)].
- [14] A. Yu. Kitaev, *Fault tolerant quantum computation by anyons*, *Annals Phys.* **303** (2003) 2 [[quant-ph/9707021](https://arxiv.org/abs/quant-ph/9707021)] [[INSPIRE](#)].

- [15] D. Radićević, *Entanglement in weakly coupled lattice gauge theories*, [arXiv:1509.08478](#) [[INSPIRE](#)].
- [16] J. Preskill, *Lecture notes on quantum computation*, <http://www.theory.caltech.edu/people/preskill/ph229/>.
- [17] C.H. Bennett, H.J. Bernstein, S. Popescu and B. Schumacher, *Concentrating partial entanglement by local operations*, *Phys. Rev. A* **53** (1996) 2046 [[quant-ph/9511030](#)] [[INSPIRE](#)].
- [18] M.A. Nielsen and I. Chuang, *Quantum computation and quantum information*, Cambridge University Press, Cambridge U.K. (2000).
- [19] M.M. Wilde, *Quantum information theory*, Cambridge University Press, Cambridge U.K. (2013) [[arXiv:1106.1445](#)].
- [20] N. Schuch, F. Verstraete and J.I. Cirac, *Nonlocal resources in the presence of superselection rules*, *Phys. Rev. Lett.* **92** (2004) 087904 [[quant-ph/0310124](#)].
- [21] N. Schuch, F. Verstraete and J.I. Cirac, *Quantum entanglement theory in the presence of superselection rules*, *Phys. Rev. A* **70** (2004) 042310 [[quant-ph/0404079](#)].
- [22] K. Van Acoleyen, N. Bultinck, J. Haegeman, M. Marien, V.B. Scholz and F. Verstraete, *The entanglement of distillation for gauge theories*, [arXiv:1511.04369](#) [[INSPIRE](#)].
- [23] V. Lahtinen, *Topological quantum computation. An analysis of an anyon model based on quantum double symmetries*, *M.Sc. thesis*, U. Helsinki, Helsinki Finland (2006).
- [24] J.-W. Chen, S.-H. Dai and J.-Y. Pang, *Strong coupling expansion of the entanglement entropy of Yang-Mills gauge theories*, [arXiv:1503.01766](#) [[INSPIRE](#)].
- [25] M. Pretko and T. Senthil, *Entanglement entropy of U(1) quantum spin liquids*, [arXiv:1510.03863](#) [[INSPIRE](#)].
- [26] J.B. Kogut and L. Susskind, *Hamiltonian formulation of Wilson's lattice gauge theories*, *Phys. Rev. D* **11** (1975) 395 [[INSPIRE](#)].