Hexadecapole strength in the rare isotopes $^{74,76}\text{Kr}$

M. Spieker^{a,*}, S.E. Agbemava^b, D. Bazin^{b,c}, S. Biswas^b, P.D. Cottle^a, P.J. Farris^{b,c},
A. Gade^{b,c}, T. Ginter^b, S. Giraud^b, K.W. Kemper^a, J. Li^b, W. Nazarewicz^{b,c}, S. Noji^b,
J. Pereira^b, L.A. Riley^d, M. Smith^b, D. Weisshaar^b, R.G.T. Zegers^{b,c}

^a Department of Physics, Florida State University, Tallahassee, FL 32306, USA

^b Facility for Rare Isotope Beams, Michigan State University, East Lansing, MI 48824, USA

^c Department of Physics and Astronomy, Michigan State University, East Lansing, MI 48824, USA

^d Department of Physics and Astronomy, Ursinus College, Collegeville, PA 19426, USA

ARTICLE INFO

Article history:

Received 17 October 2022

Received in revised form 22 February 2023

Accepted 21 April 2023

Available online 26 April 2023

Editor: B. Blank

Keywords:

Nuclear structure

Electric hexadecapole strengths

Shape coexistence

Inelastic proton scattering

Nuclear density functional theory

ABSTRACT

In the Ge-Sr mass region, isotopes with neutron number $N \leq 40$ are known to feature rapid shape changes with both nucleon number and angular momentum. To gain new insights into their structure, inelastic proton scattering experiments in inverse kinematics were performed on the rare isotopes $^{74,76}\text{Kr}$. This work focuses on observables related to the $J^\pi = 4_1^+$ states of the Kr isotopes and, in particular, on the hexadecapole degree of freedom. By performing coupled-channels calculations, hexadecapole deformation parameters β_4 were determined for the $J^\pi = 4_1^+$ states of $^{74,76}\text{Kr}$ from inelastic proton scattering cross sections. Two possible coupled-channels solutions were found. A comparison to predictions from nuclear energy density functional theory, employing both non-relativistic and relativistic functionals, clearly favors the large, positive β_4 solutions. These β_4 values are unambiguously linked to the well deformed prolate configuration. Given the $\beta_2 - \beta_4$ trend, established in this work, it appears that β_4 values could provide a sensitive measure of the nuclear shell structure.

© 2023 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

The neutron-deficient, even-even Ge, Se, Kr, and Sr isotopes exhibit rapid shape changes with both nucleon number and spin of the system [1]. Generally, the ground-state shapes appear to change from prolate to oblate towards the end of the deformed region [2–4]. Current experimental data suggest that the prolate-oblate ground-state shape transition for even-A nuclei in the Ge-Kr region occurs around neutron number $N = 36$ [5–10]. The exact location of the shape transition is still under debate and its details challenge state-of-the-art theoretical models as triaxial degrees of freedom are expected to contribute [11,12]. Additional complexity gets added as nuclei in this mass region display complex shape coexistence of oblate, prolate, spherical and triaxial configurations at low excitation energy [13,14]. Many models (see Refs. [15–19]) predict predominantly oblate-deformed ground states for nuclei around $N = 36$, with the yrast structure changing from oblate to prolate with angular momentum. This has been discussed in Ref. [8], which showed that the models that incorporate mixing between oblate and prolate configurations were able to describe both the experimentally measured $B(E2; 2_1^+ \rightarrow 0_1^+)$ and

$B(E2; 4_1^+ \rightarrow 2_1^+)$ strengths in ^{72}Kr . Interestingly, a good description of the $B(E2; 2_1^+ \rightarrow 0_1^+)$ strength in ^{72}Kr has been offered by several models while predicting different ground-state structures [8]. The 4_1^+ state and associated observables as, e.g., γ -decay probabilities or excitation cross sections could, therefore, be extremely effective discriminators between competing model descriptions.

To further investigate the sensitivity of observables associated with the 4_1^+ state to the structure of nuclei in the Ge-Sr mass region, we report on $B(E4; 4_1^+ \rightarrow 0_1^+)$ strengths and associated β_4 hexadecapole deformation parameters in $^{74,76}\text{Kr}$. These quantities were derived from inelastic proton scattering cross sections measured in inverse kinematics at the Coupled Cyclotron Facility of the National Superconducting Cyclotron Laboratory (NSCL) at Michigan State University [20] with the NSCL/Ursinus Liquid Hydrogen (LH₂) Target, the GRETINA γ -ray tracking array [21,22], and S800 magnetic spectrograph [23]. Data from low-energy Coulomb excitation (CoulEx) experiments had previously provided strong evidence for prolate ground states but also indicated significant mixing between prolate and oblate configurations [24]. Using two-state mixing arguments, a large quadrupole deformation has been attributed to the prolate configuration [24]. As will be discussed in this Letter, large intrinsic quadrupole deformations of the prolate configurations are consistent with the hexadecapole deformation

* Corresponding author.

E-mail address: mspieker@fsu.edu (M. Spieker).




Fig. 1. Doppler-corrected, in-beam γ -ray spectra for ^{76}Kr (top) and ^{74}Kr (bottom). Data are shown in black. GEANT4 simulations performed with UCGRETINA [25] are presented in red. A prompt background consisting of two exponential functions was included in the simulation. Besides transitions from the $J^\pi = 2_1^+$ and 3_1^- states, observed transitions of the populated 4^+ states are highlighted.

parameters for $^{74,76}\text{Kr}$, thus supporting the sensitivity of β_4 to the quadrupole shape of the nucleus.

For the experiments, the secondary ^{76}Kr (79% purity; ~ 4740 pps) and ^{74}Kr (51% purity; ~ 3060 pps) beams were produced from a 150 MeV/u ^{78}Kr primary beam in projectile fragmentation on a 308-mg/cm² thick ^9Be target. To select $^{74,76}\text{Kr}$ in flight, two separate A1900 magnetic settings and a 240-mg/cm² Al degrader were used. Both secondary beams were unambiguously distinguished from other components in the cocktail beam via the time-of-flight difference measured between two plastic scintillators located at the exit of the A1900 and the object position of the S800 analysis beam line. At the chosen proton center-of-mass energies of about 100 MeV, both proton and neutron matrix elements are probed [26]. The NSCL/Ursinus Liquid Hydrogen (LH₂) Target with the 8.5-mm thick cell was installed at the target position of the S800 spectrograph. The S800 magnetic spectrograph and its focal-plane detection system were used to identify the projectile-like reaction residues event-by-event from their energy loss and time of flight [23]. γ rays, which were emitted by the reaction residues in flight, were detected with the GRETINA γ -ray tracking array [21,22]. Eight GRETINA modules, containing four, 36-fold segmented HPGe detectors each, were mounted in the north half of the mounting shell to accommodate the LH₂ target. In this configuration, two modules are centered at 58°, four at 90°, and two at 122° with respect to the beam axis. At beam velocities of $v/c \approx 0.4$, event-by-event Doppler reconstruction of the residues' γ -ray energies is necessary. This reconstruction was performed based on the angle of the γ -ray emission determined from the main-interaction point and including trajectory reconstruction of the residues through the S800 spectrograph [22]. Doppler-corrected in-flight γ -ray spectra are presented in Fig. 1. The γ -ray yields were obtained by fitting γ -ray spectra, simulated with UCGRETINA [25], to the experimentally observed ones. For these fits, the ROOT [27] MINUIT2 minimizer with the default minimization algorithm MIGRAD was used [28]; as done in previous studies, see, e.g., Refs. [29–31]. Known decay branching for excited states of $^{74,76}\text{Kr}$ [32–34] was explicitly taken into account by using the GEANT4 photo-evaporation database format [35] implemented in UCGRETINA. This procedure also allowed for the correction of the γ -ray yields for observed feeders (see also Ref. [31]). Inelas-




Fig. 2. Measured kinetic-energy distributions of the outgoing $^{74,76}\text{Kr}$ beams (black) compared to results of detailed GEANT4 simulations (red) performed with UCGRETINA [25] to determine the LH₂ target thickness. Both simulations provided consistent results for the target thickness.

tic scattering cross sections were calculated from the experimental γ -ray yields by normalizing these to the number of incoming beam particles and the number of target nuclei. Pressure differences across the Kapton entrance and exit windows of the LH₂ cell cause them to bulge outwards. As first described in Ref. [29], the LH₂ target thickness was determined via a comparison of the measured kinetic-energy distribution of the reacted outgoing beam to a detailed GEANT4 simulation performed with UCGRETINA [25]. The simulation also uses the independently measured kinetic-energy distribution of the incoming beam through the empty target cell as input. The comparison is shown in Fig. 2. Excellent agreement was obtained and an areal target density of 69(3) mg/cm² was determined.

Reaction calculations were performed with the coupled-channels program CHUCK3 [36] using the optical-model parameters of [37]. For one-step processes, deformation parameters β_λ can be calculated by scaling the theoretical to the experimentally determined inelastic scattering cross sections. Excellent agreement with the adopted values [38,39] was obtained for the β_2 quadrupole deformation parameters of the 2_1^+ states, $\beta_2 = 0.40 \pm 0.02$ (stat.) ± 0.03 (sys.) for ^{76}Kr and $\beta_2 = 0.35 \pm 0.06$ (stat.) ± 0.02 (sys.) for ^{74}Kr , benchmarking both the reaction calculations and feeding correction (see also Ref. [31]). For possible multi-step processes, we followed the approach of Ref. [40] for the stable Kr isotopes. In particular, coupled-channels calculations were performed to determine the hexadecapole deformation parameter, β_4 , from the inelastic scattering cross sections measured for the 4_1^+ states of $^{74,76}\text{Kr}$. The inelastic scattering cross sections to the 4_1^+ state are 5.1(5) mb for ^{76}Kr and 6.1(11) mb for ^{74}Kr , respectively. Additional feeding contributions coming from unobserved feeders cannot be ruled out entirely. For the two-step contribution via the intermediate 2_1^+ state, the adopted β_2 values were chosen [38,39]. In principle, this leaves β_4 as the only free parameter to match the experimental cross section. Using this approach, values of $\beta_4 = 0.201 \pm 0.009$ (stat.) ± 0.016 (sys.) for ^{76}Kr and $\beta_4 = 0.23 \pm 0.02$ (stat.) ± 0.02 (sys.) for ^{74}Kr were determined. Systematic uncertainties include the uncertainty of the adopted β_2 value [38,39] and uncertainties in the reaction kinematics due to the target thickness. Systematic uncertainties due to unobserved feeders cannot be estimated reliably. It is, however, important to state that if there existed a single feeder or even several feeders leading to a further feeding-subtracted β_4 value of around +0.1 as in ^{78}Kr [40], then we should have been able to observe the corresponding γ -ray transitions up to an energy of at least 2 MeV. We also note that there exist alternative coupled-channels solutions corresponding to negative β_4 values. These values are $\beta_4 = -0.127 \pm 0.009$ (stat.) ± 0.022 (sys.) for ^{76}Kr and

$\beta_4 = -0.17 \pm 0.02$ (stat.) ± 0.02 (sys.). Positive and negative β_4 solutions from inelastic scattering cross sections were also reported in the rare-earth region [41]; they correspond to constructive and destructive interference between one-step and two-step processes. In this context, we want to stress that two-step excitation alone would only provide cross sections of ~ 0.8 mb for ^{76}Kr and ~ 0.5 mb for ^{74}Kr , respectively. A significant direct (one-step) contribution is therefore needed to explain the experimental data. As we will discuss later, the positive solutions for β_4 are preferred. We note, however, that the measured inelastic scattering cross sections have no sensitivity to the sign of β_4 . Therefore, both values, which were determined through the coupled-channels analysis, are reported in this Letter.

The β_2 deformation parameters determined in our experiments [31], the adopted values [38,39], as well as the calculated β_2 values for the unperturbed prolate configuration are shown in Fig. 3 (a). The latter were determined by Clément *et al.* through a two-state mixing calculation as described in Ref. [24]. The positive β_4 deformation parameters for $^{74,76}\text{Kr}$ are shown in Fig. 3 (b), which also includes the data of Ref. [40] for isotopes with $A \geq 78$. A significant increase of β_4 is observed in $^{74,76}\text{Kr}$. Fig. 3 (c) shows the evolution of the $B(E4; 4_1^+ \rightarrow 0_1^+)$ strength along the Kr isotopic chain. Consistent with Ref. [40], the $B(E4)$ strengths for $^{74,76}\text{Kr}$ were calculated from the β_4 parameters by assuming a uniform mass distribution of radius $R = 1.2 \text{ fm} \times A^{1/3}$ [42]. Using the positive β_4 values, significant $B(E4; 4_1^+ \rightarrow 0_1^+)$ strengths of 22.7 ± 1.0 (stat.) ± 1.8 (sys.) W.u. for ^{76}Kr and 29 ± 2 (stat.) ± 2 (sys.) W.u. for ^{74}Kr are determined. The $B(E4)$ strengths, determined from the negative β_4 solutions, are also shown in Fig. 3 (b). The $B(E4; 4_1^+ \rightarrow 0_1^+)$ strengths (and β_4 values) of the corresponding Se and Ge isotones are significantly smaller [43,44]. It is worth noting that strengths larger than 10 W.u. were also observed in the $A = 100$ region [45] as well as in the Nd isotopes [46] in inelastic light-ion scattering experiments. Besides the $J^\pi = 4_1^+$ state of ^{76}Kr , the 1957-keV, $J^\pi = 4_2^+$ state was also populated in our (p, p') experiment (see Fig. 1). As this state does not belong to the ground-state band of ^{76}Kr , single-step excitation was assumed and $\beta_4 = 0.151 \pm 0.011$ (stat.) ± 0.012 (sys.) derived. This β_4 value corresponds to a $B(E4)$ strength of 12.9 ± 0.9 (stat.) ± 2.0 (sys.) W.u., which brings the total $B(E4)$ strength up to 36 ± 2 (stat.) ± 4 (sys.) W.u. in ^{76}Kr . No higher-lying, excited 4_1^+ states were observed for ^{74}Kr . In the following, we concentrate on the $J^\pi = 4_1^+$ states of the Kr isotopes and associated observables.

To investigate the origin of the large β_4 values in $^{74,76}\text{Kr}$, nuclear density functional theory (DFT) calculations were performed using the Skyrme SkM* [47] and UNEDF1 [48] energy density functionals, as well as the covariant NL3* [49] and DD-PC1 [50] energy density functionals. As in Ref. [51], the mixed-type density-dependent delta interaction [52] with the Lipkin-Nogami approximate particle-number projection was used in the pairing channel for the Skyrme functionals (SEDFs). The separable finite-range pairing [53] with the strength defined as in Ref. [54] was adopted for the covariant functionals (CDFTs). For the SEDFs, calculations were performed with the parallel DFT axial solver HFBTHO [55]. For the CDFTs, the calculations were carried out employing an axial solver of Ref. [54].

The charge multipole moments $Q_{\lambda 0}$ ($\lambda = 2, 4$) and dimensionless deformation parameters β_λ are defined as:

$$Q_{\lambda 0} = \sqrt{\frac{16\pi}{2\lambda + 1}} \langle r^\lambda Y_{\lambda 0} \rangle = \frac{3}{\sqrt{(2\lambda + 1)\pi}} Z R_0^\lambda \beta_\lambda, \quad (1)$$

where $R_0 = 1.2A^{1/3}$. For strongly deformed nuclei, deformation parameters β_λ defined through the linear relation (1) can strongly deviate from commonly used deformation parameters entering the




Fig. 3. (a) Experimental β_2 values [new data (black circles), adopted values [38,39] (gray circles)], and values determined by Clément *et al.* for the unperturbed prolate configuration [24] (white squares)], (b) β_4 values [new data (black diamonds) and previous (p, p') data [40] (gray diamonds)], and (c) experimental $B(E4; 4_1^+ \rightarrow 0_1^+)$ strengths in neutron-deficient Kr isotopes. Alternative values, obtained for the negative β_4 values discussed in the text, are presented with white diamonds in panel (c). Theoretical DFT predictions for the corresponding observables are shown in panels (a) and (b). These were obtained using the energy density functionals SkM* (black longer dashed line) and UNEDF1 (red solid line), as well as two covariant energy density functionals DD-PC1 (orange dotted line) and NL3* (blue short dashed line). For β_2 , predictions are shown for the prolate minimum only. In addition, predictions for the oblate minimum made with SkM* are shown (green dotted-dashed line) in (b).

multipole expansion of the nuclear surface [2,56]. Consequently, one has to exercise caution when comparing to experimental data.

The predicted trend of β_4 with mass number A is shown in Fig. 3(b). Keeping the linear approximation of Eq. (1) in mind, all our models predict a significant increase in β_4 for the prolate minima of $^{74,76,78}\text{Kr}$. The results obtained with the functional SkM*, which predicts prolate ground states with large quadrupole deformation $\beta_2 = 0.46 - 0.49$ for $^{74,76}\text{Kr}$ [see Fig. 3(a)], most closely resemble the observed experimental trend for the positive β_4 coupled-channels solution; also in absolute magnitude [see Fig. 3(b)]. In fact, Clément *et al.* calculated a β_2 value of ~ 0.5 for the unperturbed prolate configuration in $^{74,76}\text{Kr}$ using two-state mixing with an unperturbed oblate configuration [24]. This value is in excellent agreement with our DFT predictions for the prolate minimum [see Fig. 3(a)]. We note that UNEDF1, DD-PC1, and NL3* yield an oblate ground state for ^{74}Kr . In conflict with both coupled-channels solutions, small β_4 values of ~ -0.02 are predicted at the oblate configurations. The predictions made with SkM* for the oblate minimum are shown in Fig. 3 (b). They do not describe the experimentally observed trend emphasizing that large, positive β_4 values are associated with the prolate minima.

To show the connection between the large β_2 and β_4 values, both the experimentally determined and theoretically predicted deformations are compiled in Fig. 4. It is seen that large values of β_4 correspond to large values of β_2 both in experiment and theory. In fact, non-relativistic (SkM* and UNEDF1) and relativistic (DD-PC1 and NL3*) functionals conform to a remarkably similar trend [see Fig. 4 (b)]. For nuclei with $\beta_2 \geq 0.3$, excellent agreement is observed between theory and experiment. The deviation of the experimental data for $^{74,76}\text{Kr}$ from this trend, when using the adopted β_2 values [38,39], further supports the shape-mixing




Fig. 4. Hexadecapole deformation parameter β_4 plotted against β_2 : (a) experimental values; (b) theoretical predictions. Earlier data for β_4 are from Ref. [40] (gray circles). The experimental β_2 values were taken from Ref. [38,39]. The open circles in (a) correspond to the β_2 values calculated from the quadrupole moments Q_{20} of the unperturbed prolate configurations reported in Ref. [24]. The SkM* predictions are marked in panel (a) with a dashed line.

hypothesis of Ref. [24]. Considering that the β_2 values are smaller than 0.3 in the corresponding Ge and Se isotones [38,39], this connection also naturally explains why hexadecapole deformations are significantly smaller [43,44]. For instance, the $N = 40$ isotones of ^{76}Kr , ^{72}Ge [$\beta_2 = 0.240(2)$] and ^{74}Se [$\beta_2 = 0.290(8)$] have small β_4 values below 0.05.

A simple interpretation of the positive hexadecapole moments can be obtained by using the geometric idea of the polar-gap model [57] developed to explain β_4 deformations of rare-earth nuclei [58]. According to this model, positive deformations β_4 are expected at the beginning of a shell and negative hexadecapole deformations are expected at the end of a shell (see also Refs. [59–64]). The collective properties of $A = 70 - 80$ nuclei are governed by deformed orbitals originating from the $0g_{9/2}$ unique-parity shell [65]. At large prolate quadrupole deformations $\beta_2 = 0.4 - 0.5$, the Nilsson levels [440]1/2 and [431]3/2 at the bottom of the $0g_{9/2}$ shell become occupied and this puts $^{74,76}\text{Kr}$ right at the beginning of the deformed shell with the low- K Nilsson orbitals being at the Fermi surface [65]. As also discussed in Ref. [64], these low- K orbitals are primarily concentrated in the equatorial plane of the quadrupole deformed core; hence, their population contributes to a positive hexadecapole moment.

In summary, we have performed inelastic proton scattering experiments in inverse kinematics on the rare isotopes $^{74,76}\text{Kr}$ with GREINA, the S800 spectrograph and NSCL/Ursinus LH_2 target. In this Letter, we report $B(E4)$ strengths and β_4 deformations of these nuclei. Based on a coupled-channels analysis, two possible solutions for the β_4 hexadecapole deformation parameters of the $J^\pi = 4_1^+$ states in $^{74,76}\text{Kr}$ were determined, differing in sign because of the interference between one-step and two-step excitations. However, the two-step contributions are weak. Nuclear DFT calculations, employing both non-relativistic and relativistic energy density functionals, strongly favor the solutions with large positive values of β_4 . For this scenario, very good agreement between experiment and theory was obtained. This finding supports the results of Ref. [24] suggesting predominantly prolate structures of the $^{74,76}\text{Kr}$ ground-, $J^\pi = 2_1^+$, and $J^\pi = 4_1^+$ states. The large positive β_4 values reported in this Letter are unambiguously linked to the prolate configurations. Given the remarkable correlation between β_2 and β_4 values predicted by our models for the Kr isotopes, it suggests that β_4 deformations could indeed be a sensitive

indicator of prolate shapes. Since the prolate-oblate ground-state phase transition presumably happens at $A = 72$ (or $N = 36$), it is a valid question what one might observe for the hexadecapole moments of $^{70,72}\text{Kr}$ and other nuclei in the neutron-deficient Ge-Sr region. Future experiments at rare isotope beam facilities will help answering this question.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

Data will be made available on request.

Acknowledgements

This work was supported by the National Science Foundation (NSF) under Grant No. PHY-2012522 (WoU-MMA: Studies of Nuclear Structure and Nuclear Astrophysics), Grant No. PHY-1565546 (NSCL), Grant No. PHY-2209429 (Windows on the Universe: Nuclear Astrophysics at FRIB), by the Department of Energy, Office of Science, Office of Nuclear Physics, Grant Nos. DE-SC0020451 and DE-SC0013365 (MSU), and by the Department of Energy, NNSA, Grant No. DOE-DE-NA0004074 (the Stewardship Science Academic Alliances program). GREINA was funded by the Department of Energy, Office of Science. The operation of the array at NSCL was supported by the Department of Energy under Grant No. DE-SC0019034. M.S. acknowledges support through the FRIB Visiting Scholar Program for Experimental Science 2020.

References

- [1] J. Eberth, R.A. Meyer, K. Sistemich (Eds.), *Nuclear Structure of the Zirconium Region*, Springer, Berlin Heidelberg, 1988.
- [2] W. Nazarewicz, I. Ragnarsson, Nuclear deformations, in: D. Poenaru, W. Greiner (Eds.), *Handbook of Nuclear Properties*, Clarendon Press, 1996, p. 80.
- [3] J. Jolie, A. Linnemann, *Phys. Rev. C* 68 (2003) 031301.
- [4] X.Q. Yang, L.J. Wang, J. Xiang, X.Y. Wu, Z.P. Li, *Phys. Rev. C* 103 (2021) 054321.
- [5] R. Lecomte, M. Irshad, S. Landsberger, P. Paradis, S. Monaro, *Phys. Rev. C* 22 (1980) 1530.
- [6] A. Gade, D. Bazin, A. Becerril, C.M. Campbell, J.M. Cook, D.J. Dean, D.-C. Dinca, T. Glasmacher, G.W. Hitt, M.E. Howard, W.F. Mueller, H. Olliver, J.R. Terry, K. Yoneda, *Phys. Rev. Lett.* 95 (2005) 022502.
- [7] J. Jungvall, A. Görgen, M. Girod, J.-P. Delaroche, A. Dewald, C. Dossat, E. Farnea, W. Korten, B. Melon, R. Menegazzo, A. Obertelli, R. Orlandi, P. Petkov, T. Pissulla, S. Siem, R.P. Singh, J. Srebrny, C. Theisen, C.A. Ur, J.J. Valiente-Dobón, K.O. Zell, M. Zielińska, *Phys. Rev. Lett.* 100 (2008) 102502.
- [8] H. Iwasaki, A. Lemasson, C. Morse, A. Dewald, T. Braunroth, V.M. Bader, T. Baugher, D. Bazin, J.S. Berryman, C.M. Campbell, A. Gade, C. Langer, I.Y. Lee, C. Loelius, E. Lunderberg, F. Recchia, D. Smalley, S.R. Stroberg, R. Wadsworth, C. Walz, D. Weisshaar, A. Westerberg, K. Whitmore, K. Wimmer, *Phys. Rev. Lett.* 112 (2014) 142502.
- [9] J. Henderson, C.Y. Wu, J. Ash, P.C. Bender, B. Elman, A. Gade, M. Grindler, H. Iwasaki, E. Kwan, B. Longfellow, T. Mijatović, D. Rhodes, M. Spieker, D. Weisshaar, *Phys. Rev. Lett.* 121 (2018) 082502.
- [10] K. Wimmer, T. Arici, W. Korten, P. Doornenbal, J.-P. Delaroche, M. Girod, J. Libert, T.R. Rodríguez, P. Aguilera, A. Algorta, T. Ando, H. Baba, B. Blank, A. Boso, S. Chen, A. Corsi, P. Davies, G. de Angelis, G. de France, D.T. Doherty, J. Gerl, R. Gernhäuser, T. Goigoux, D. Jenkins, G. Kiss, S. Koyama, T. Motobayashi, S. Nagamine, M. Niikura, S. Nishimura, A. Obertelli, D. Lubos, V.H. Phong, B. Rubio, E. Sahin, T.Y. Saito, H. Sakurai, L. Sinclair, D. Steppenbeck, R. Taniuchi, V. Vaquero, R. Wadsworth, J. Wu, M. Zielińska, *Eur. Phys. J. A* 56 (2020) 159.
- [11] A.D. Ayangeakaa, R.V.F. Janssens, C.Y. Wu, J.M. Allmond, J.L. Wood, S. Zhu, M. Albers, S. Almaraz-Calderon, B. Bucher, M.P. Carpenter, C.J. Chiara, D. Cline, H.L. Crawford, H.M. David, J. Harker, A.B. Hayes, C.R. Hoffman, B.P. Kay, K. Kolos, A. Korichi, T. Lauritsen, A.O. Macchiavelli, A. Richard, D. Seweryniak, A. Wiens, *Phys. Lett. B* 754 (2016) 254.
- [12] J. Henderson, C.Y. Wu, J. Ash, B.A. Brown, P.C. Bender, R. Elder, B. Elman, A. Gade, M. Grindler, H. Iwasaki, B. Longfellow, T. Mijatović, D. Rhodes, M. Spieker, D. Weisshaar, *Phys. Rev. C* 99 (2019) 054313.

- [13] K. Heyde, J.L. Wood, *Rev. Mod. Phys.* **83** (2011) 1467.
- [14] P.E. Garrett, M. Zielińska, E. Clément, *Prog. Part. Nucl. Phys.* **124** (2022) 103931.
- [15] M. Bender, P. Bonche, P.-H. Heenen, *Phys. Rev. C* **74** (2006) 024312.
- [16] M. Girod, J.-P. Delaroche, A. Görgen, A. Obertelli, *Phys. Lett. B* **676** (2009) 39.
- [17] N. Hinohara, K. Sato, T. Nakatsukasa, M. Matsuo, K. Matsuyanagi, *Phys. Rev. C* **82** (2010) 064313.
- [18] K. Sato, N. Hinohara, *Nucl. Phys. A* **849** (2011) 53.
- [19] T.R. Rodríguez, *Phys. Rev. C* **90** (2014) 034306.
- [20] A. Gade, B.M. Sherrill, *Phys. Scr.* **91** (2016) 053003.
- [21] S. Paschalis, I.Y. Lee, A.O. Macchiavelli, C.M. Campbell, M. Cromaz, S. Gros, J. Pavan, J. Qian, R.M. Clark, H.L. Crawford, D. Doering, P. Fallon, C. Lionberger, T. Loew, M. Petri, T. Stezelberger, S. Zimmermann, D.C. Radford, K. Lagergren, D. Weisshaar, R. Winkler, T. Glasmacher, J.T. Anderson, C.W. Beausang, *Nucl. Instrum. Methods Phys. Res., Sect. A* **709** (2013) 44.
- [22] D. Weisshaar, D. Bazin, P.C. Bender, C.M. Campbell, F. Recchia, V. Bader, T. Baugher, J. Belarge, M.P. Carpenter, H.L. Crawford, M. Cromaz, B. Elman, P. Fallon, A. Forney, A. Gade, J. Harker, N. Kobayashi, C. Langer, T. Lauritsen, I.Y. Lee, A. Lemasson, B. Longfellow, E. Lunderberg, A.O. Macchiavelli, K. Miki, S. Momiyama, S. Noji, D.C. Radford, M. Scott, J. Sethi, S.R. Stroberg, C. Sullivan, R. Titus, A. Wiens, S. Williams, K. Wimmer, S. Zhu, *Nucl. Instrum. Methods Phys. Res., Sect. A* **847** (2017) 187.
- [23] D. Bazin, J.A. Caggiano, B.M. Sherrill, J. Yurkon, A. Zeller, *Nucl. Instrum. Methods Phys. Res. B* **204** (2003) 629.
- [24] E. Clément, A. Görgen, W. Kortén, E. Bouchez, A. Chatillon, J.-P. Delaroche, M. Girod, H. Goutte, A. Hürstel, Y.L. Coz, A. Obertelli, S. Péru, C. Theisen, J.N. Wilson, M. Zielińska, C. Andreoiu, F. Becker, P.A. Butler, J.M. Casandjian, W.N. Catford, T. Czornyka, G.d. France, J. Gerl, R.-D. Herzberg, J. Iwanicki, D.G. Jenkins, G.D. Jones, P.J. Napiorkowski, G. Sletten, C.N. Timis, *Phys. Rev. C* **75** (2007) 054313.
- [25] L.A. Riley, D. Weisshaar, H.L. Crawford, M.L. Agiorgousis, C.M. Campbell, M. Cromaz, P. Fallon, A. Gade, S.D. Gregory, E.B. Haldeman, L.R. Jarvis, E.D. Lawson-John, B. Roberts, B.V. Sadler, C.G. Stine, *Nucl. Instrum. Methods Phys. Res., Sect. A* **1003** (2021) 165305.
- [26] P. Cottle, *Nucl. Phys. A* **682** (2001) 124.
- [27] R. Brun, F. Rademakers, New computing techniques in physics research V, *Nucl. Instrum. Methods Phys. Res., Sect. A, Accel. Spectrom. Detect. Assoc. Equip.* **389** (81) (1997).
- [28] Minuit2 User Guide, <https://root.cern.ch/root/html/doc/guides/minuit2/Minuit2.html>, 2023.
- [29] L.A. Riley, D. Bazin, J. Belarge, P.C. Bender, B.A. Brown, P.D. Cottle, B. Elman, A. Gade, S.D. Gregory, E.B. Haldeman, K.W. Kemper, B.R. Klybor, M.A. Liggett, S. Lipschutz, B. Longfellow, E. Lunderberg, T. Mijatovic, J. Pereira, L.M. Skiles, R. Titus, A. Volya, D. Weisshaar, J.C. Zamora, R.G.T. Zegers, *Phys. Rev. C* **100** (2019) 044312.
- [30] A.M. Hill, A. Gade, D. Bazin, B.A. Brown, B. Elman, P. Farris, J. Li, B. Longfellow, J. Pereira, A. Revel, D. Rhodes, M. Spieker, D. Weisshaar, *Phys. Rev. C* **104** (2021) 014305.
- [31] M. Spieker, L.A. Riley, P.D. Cottle, K.W. Kemper, D. Bazin, S. Biswas, P.J. Farris, A. Gade, T. Ginter, S. Giraud, J. Li, S. Noji, J. Pereira, M. Smith, D. Weisshaar, R.G.T. Zegers, *Phys. Rev. C* **106** (2022) 054305.
- [32] ENSDF, NNDC Online Data Service, ENSDF database, <http://www.nndc.bnl.gov/ensdf/>, 2023.
- [33] A. Giannatiempo, A. Perego, P. Sona, A. Nannini, H. Mach, B. Fogelberg, M.J.G. Borge, O. Tengblad, L.M. Fraile, A.J. Aas, K. Gulda, *Phys. Rev. C* **72** (2005) 044308.
- [34] R. Dunlop, G.C. Ball, J.R. Leslie, C.E. Svensson, I.S. Towner, C. Andreoiu, S. Chagnon-Lessard, A. Chester, D.S. Cross, P. Finlay, A.B. Garnsworthy, P.E. Garrett, J. Glistler, G. Hackman, B. Hadinia, K.G. Leach, E.T. Rand, K. Starosta, E.R. Tardiff, S. Triambak, S.J. Williams, J. Wong, S.W. Yates, E.F. Zganjar, *Phys. Rev. C* **88** (2013) 045501.
- [35] S. Agostinelli, J. Allison, K. Amako, J. Apostolakis, H. Araujo, P. Arce, M. Asai, D. Axen, S. Banerjee, G. Barrand, F. Behner, L. Bellagamba, J. Boudreau, L. Broglia, A. Brunengo, H. Burkhardt, S. Chauvie, J. Chuma, R. Chytracsek, G. Cooperman, G. Cosmo, P. Degtyarenko, A. Dell'Acqua, G. Depaola, D. Dietrich, R. Enami, A. Feliciello, C. Ferguson, H. Fesefeldt, G. Folger, F. Foppiano, A. Forti, S. Garelli, S. Giani, R. Giannitrapani, D. Gibin, J. Gómez Cadenas, I. González, G. Gracia Abril, G. Greeniaus, W. Greiner, V. Grichine, A. Grossheim, S. Guatelli, P. Gumplinger, R. Hamatsu, K. Hashimoto, H. Hasui, A. Heikkinen, A. Howard, V. Ivanchenko, A. Johnson, F. Jones, J. Kallenbach, N. Kanaya, M. Kawabata, Y. Kawabata, M. Kawaguti, S. Kelner, P. Kent, A. Kimura, T. Kodama, R. Kokoulin, M. Kossov, H. Kurashige, E. Lamanna, T. Lampén, V. Lara, V. Lefebvre, F. Lei, M. Liendl, W. Lockman, F. Longo, S. Magni, M. Maire, E. Medernach, K. Minamimoto, P. Mora de Freitas, Y. Morita, K. Murakami, M. Nagamatsu, R. Nartallo, P. Nieminen, T. Nishimura, K. Ohtsubo, M. Okamura, S. O'Neale, Y. Oohata, K. Paech, J. Perl, A. Pfeiffer, M. Pia, F. Ranjard, A. Rybin, S. Sadilov, E. Di Salvo, G. Santin, T. Sasaki, N. Savvas, Y. Sawada, S. Scherer, S. Sei, V. Sirotenko, D. Smith, N. Starkov, H. Stoecker, J. Sulkimo, M. Takahata, S. Tanaka, E. Tcherniaev, E. Safai Tehrani, M. Tropeano, P. Truscott, H. Uno, L. Urban, P. Urban, M. Verderi, A. Walkden, W. Wander, H. Weber, J. Wellisch, T. Wenaus, D. Williams, D. Wright, T. Yamada, H. Yoshida, D. Zschesche, *Nucl. Instrum. Methods Phys. Res., Sect. A, Accel. Spectrom. Detect. Assoc. Equip.* **506** (2003) 250.
- [36] P.D. Kunz, J.R. Comfort, Program CHUCK3 (1984), unpublished.
- [37] A.J. Koning, J.P. Delaroche, *Nucl. Phys. A* **713** (2003) 231.
- [38] B. Pritychenko, M. Birch, M. Horoi, B. Singh, *Nucl. Data Sheets* **120** (2014) 112.
- [39] B. Pritychenko, M. Birch, B. Singh, M. Horoi, *At. Data Nucl. Data Tables* **107** (2016) 1.
- [40] N. Sakamoto, S. Matsuki, K. Ogino, Y. Kadota, T. Tanabe, Y. Okuma, *Phys. Lett. B* **83** (1979) 39.
- [41] R.M. Ronningen, J.H. Hamilton, L. Varnell, J. Lange, A.V. Ramayya, G. Garcia-Bermudez, W. Lourens, L.L. Riedinger, F.K. McGowan, P.H. Stelson, R.L. Robinson, J.L.C. Ford, *Phys. Rev. C* **16** (1977) 2208.
- [42] A.M. Bernstein, Isoscalar transition rates in nuclei from the (α, α') reaction, in: M. Baranger, E. Vogt (Eds.), *Advances in Nucl. Phys.*, Springer US, Boston, MA, 1969, pp. 325–476.
- [43] K. Ogino, *Phys. Rev. C* **33** (1986) 71.
- [44] B. Schürmann, D. Rychel, B. van Kruchten, J. Speer, C.A. Wiedner, *Nucl. Phys. A* **475** (1987) 361.
- [45] M. Pignatelli, N. Blasi, S. Micheletti, R. De Leo, L. LaGamba, R. Perrino, J. Bordewijk, M. Hofstee, J. Schippers, S. van der Werf, J. Wesseling, M. Harakeh, *Nucl. Phys. A* **540** (1992) 27.
- [46] M. Pignatelli, N. Blasi, J. Bordewijk, R. De Leo, M. Harakeh, M. Hofstee, S. Micheletti, R. Perrino, V. Ponomarev, V. Soloviev, A. Sushkov, S. van der Werf, *Nucl. Phys. A* **559** (1993) 1.
- [47] J. Bartel, P. Quentin, M. Brack, C. Guet, H.-B. Håkansson, *Nucl. Phys. A* **386** (1982) 79.
- [48] M. Kortelainen, J. McDonnell, W. Nazarewicz, P.-G. Reinhard, J. Sarich, N. Schunck, M.V. Stoitsov, S.M. Wild, *Phys. Rev. C* **85** (2012) 024304.
- [49] G. Lalazisis, S. Karatzikos, R. Fossion, D.P. Arteaga, A. Afanasjev, P. Ring, *Phys. Lett. B* **671** (2009) 36.
- [50] T. Nikšić, D. Vretenar, P. Ring, *Phys. Rev. C* **78** (2008) 034318.
- [51] M.V. Stoitsov, J. Dobaczewski, W. Nazarewicz, S. Pittel, D.J. Dean, *Phys. Rev. C* **68** (2003) 054312.
- [52] J. Dobaczewski, W. Nazarewicz, M.V. Stoitsov, *Eur. Phys. J. A* **15** (2002) 21.
- [53] Y. Tian, Z. Ma, P. Ring, *Phys. Lett. B* **676** (2009) 44.
- [54] S.E. Agbemava, A.V. Afanasjev, D. Ray, P. Ring, *Phys. Rev. C* **89** (2014) 054320.
- [55] R.N. Perez, N. Schunck, R.-D. Lasserri, C. Zhang, J. Sarich, *Comput. Phys. Commun.* **220** (2017) 363.
- [56] G.A. Leander, Y.S. Chen, *Phys. Rev. C* **37** (1988) 2744.
- [57] G. Bertsch, *Phys. Lett. B* **26** (1968) 130.
- [58] D. Hendrie, N. Glendenning, B. Harvey, O. Jarvis, H. Duhm, J. Saudinos, J. Mahoney, *Phys. Lett. B* **26** (1968) 127.
- [59] J. Jänecke, *Phys. Lett. B* **103** (1981) 1.
- [60] P. Möller, A. Sierk, T. Ichikawa, H. Sagawa, *At. Data Nucl. Data Tables* **109–110** (2016) 1.
- [61] W. Nazarewicz, P. Rozmej, *Nucl. Phys. A* **369** (1981) 396.
- [62] T. Ichihara, H. Sakaguchi, M. Nakamura, M. Yosoi, M. Ieiri, Y. Takeuchi, H. Togawa, T. Tsutsumi, S. Kobayashi, *Phys. Lett. B* **182** (1986) 301.
- [63] T. Ichihara, H. Sakaguchi, M. Nakamura, M. Yosoi, M. Ieiri, Y. Takeuchi, H. Togawa, T. Tsutsumi, S. Kobayashi, *Phys. Rev. C* **36** (1987) 1754.
- [64] R.F. Casten, Nuclear structure from a simple perspective, in: P.E. Hodgson (Ed.), *Oxford Studies in Nucl. Phys.*, Oxford Science Publications, NY, USA, 2000, pp. 381–385.
- [65] W. Nazarewicz, J. Dudek, R. Bengtsson, T. Bengtsson, I. Ragnarsson, *Nucl. Phys. A* **435** (1985) 397.