



# Virasoro symmetries of the constrained dispersionless mKP hierarchy

Xiaoman Luo, Chuanzhong Li\*

College of Mathematics and Systems Science, Shandong University of Science and Technology, Qingdao, 266590, Shandong, PR China

## ARTICLE INFO

### Article history:

Received 18 April 2023

Received in revised form 25 June 2023

Accepted 28 June 2023

Available online 3 July 2023

Editor: N. Lambert

### Keywords:

Constrained dispersionless mKP hierarchy

Auto-Bäcklund transformations

Dispersionless Miura map

Additional symmetries

Virasoro symmetries

## ABSTRACT

This paper aims at the auto-Bäcklund transformations for the dispersionless mKP hierarchy, dispersionless Miura map between the constrained dispersionless mKP hierarchy and the constrained dispersionless Harry Dym hierarchy and Virasoro symmetries of the constrained dispersionless mKP hierarchy. The additional symmetries of the constrained dispersionless mKP hierarchy are constructed in a manner similar to that of the dispersionless mKP hierarchy. By comparison, a new Laurent series  $\mathcal{Y}_l$  is introduced. The additional flows form a subalgebra of the Virasoro algebra.

© 2023 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP<sup>3</sup>.

## 1. Introduction

The Kadomtsev-Petviashvili (KP) hierarchy is one of the most important integrable hierarchies and it appears in many different fields of mathematics and physics, such as the enumerative algebraic geometry, topological field and string theory [1]. The KP hierarchy has two important symmetries, they are the Sato's Bäcklund symmetry generated by vertex operator and the additional symmetries by Orlov-Schulman operator [2]. The KP hierarchy has many generalizations and extensions including universal character hierarchy [3–8]. The mKP hierarchy is a modified KP hierarchy. As one of the most important research objects in mathematical physics and integrable systems, the mKP hierarchy is of great significance [9]. Dispersionless integrable hierarchies (KP, Toda etc.) are important systems in integrable field theories, especially applied in mathematics, physics, topological field theory and matrix model theory [10]. We introduce the Planck constant  $\epsilon$  into the ordinary KP hierarchy, and the Planck constant  $\epsilon$  is restricted to approach 0. The order of microdifferential operators with  $\epsilon$  are defined by

$$\text{ord}^\epsilon(\sum a_{l,j}(t)\epsilon^l\partial^j) = \max\{j - l | a_{l,j}(t) \neq 0\}.$$

In particular,  $\text{ord}^\epsilon(\epsilon) = -1$ ,  $\text{ord}^\epsilon(\partial) = 1$ . The principal symbol of a microdifferential operator  $A = \sum a_{l,j}(t)\epsilon^l\partial^j$  is

$$\sigma^\epsilon(A) = \epsilon^{-\text{ord}^\epsilon(A)} \sum_{j-l=\text{ord}^\epsilon(A)} a_{l,j}k^j,$$

and Lie bracket has become Poisson bracket.

Gauge transformation is one kind of powerful methods to construct the solutions of the integrable systems, it has significant significance and is widely used [11–17]. For soliton theory, the Miura map has played a significant role in its development [18–20]. The Miura map is generally not easy to solve and is a transformation between two nonlinear evolution equations [21,22]. However, the solution of one nonlinear system can be obtained from the solution of the other nonlinear system through the Miura map. Bäcklund transformations can be used to study the soliton hierarchies, which is a very effective tool. On the basis of appropriate gauge transformation, using commonly used dressing method can obtain the  $(2+1)$ -dimensional transformations [23]. In [19], the Miura map between the dispersionless KP hierarchy and dispersionless modified KP hierarchy is given. This article has constructed

\* Corresponding author.

E-mail address: [lichuanzhong@sdust.edu.cn](mailto:lichuanzhong@sdust.edu.cn) (C. Li).

$$\mathcal{L}_{dmKP} = e^{-ad\phi} \mathcal{L}_{dKP},$$

which can also be used for the Miura map between the constrained dispersionless KP hierarchy and constrained dispersionless modified KP hierarchy and the dKP hierarchy with self-consistent sources to the dmKP hierarchy with self-consistent sources [23]. In [20], the Miura map between the dDym hierarchy and the dmKP hierarchy is given. Its structure can also be used for the Miura map between the constrained dispersionless Harry Dym hierarchy and constrained dispersionless modified KP hierarchy [22]. The constrained dispersionless dmKP hierarchy can be thought of as the quasi-classical limit of the constrained mKP hierarchy [24].

Additional symmetries are important symmetries that explicitly depend on time and space [25–27]. Additional symmetries do not commute with each other and the additional symmetry flows of the KP hierarchy can form an infinite-dimensional algebra  $W_{1+\infty}$ . Additional symmetry also has wide applications [28,29]. In [30], the additional symmetries of the dispersionless cKP hierarchy are given by introducing a new Laurent series  $\mathcal{Y}_l$ . Additional symmetry can be used to derive string equations and Virasoro constraints in string theory [31,32].

This paper is organized in the following way. In Section 2, some basic knowledge will be introduced. We introduce the mKP hierarchy, the constrained mKP hierarchy, the dmKP hierarchy, the constraints of dispersionless mKP hierarchy, the Harry Dym hierarchy, the dispersionless Harry Dym hierarchy and the constrained dispersionless Harry Dym hierarchy. In Section 3, we give the auto-Bäcklund transformations for dmKP hierarchy. In Section 4, we give dispersionless Miura map between the constrained dispersionless mKP hierarchy and the constrained dispersionless Harry Dym hierarchy. In Section 5, the additional symmetries of the constrained dispersionless mKP hierarchy are given. In Section 6, the conclusions and discussions are given.

## 2. Background materials

In section 2, we introduce some basic knowledge, the mKP hierarchy, the constrained mKP hierarchy, the dmKP hierarchy, the constrained dispersionless mKP hierarchy, the Harry Dym hierarchy, the dispersionless Harry Dym hierarchy and the constrained dispersionless Harry Dym hierarchy, which we will use them later on.

The Lax operator of the mKP hierarchy is as

$$L_{mKP} = \partial + v_0 + v_1 \partial^{-1} + v_2 \partial^{-2} + \dots,$$

where  $\partial = \partial_x, v_i = v_i(x = t_1, t_2, \dots)$ .

And it satisfies the Lax equations

$$\partial_{t_k} L_{mKP} = [Q_k, L_{mKP}], \tag{2.1}$$

where  $Q_k = (L_{mKP}^k)_{\geq 1}$  means the part of order  $\geq 1$  of  $L_{mKP}^k$ .

The Lax equation (2.1) is equivalent to the existence of wave function  $\Psi(t, \lambda)$  such that  $(t = (t_1 = x, t_2, \dots))$

$$L_{mKP} \Psi = \lambda \Psi,$$

$$\partial_{t_k} \Psi = Q_k \Psi,$$

where  $\lambda$  is the eigenvalue of  $L_{mKP}$ .

Let  $p = \sum_{i=-\infty}^m p_i \partial^i$  be an arbitrary pseudo-differential operators,  $p^* = \sum_{i=-\infty}^m (-1)^i \partial^i p_i$  is its formal adjoint operator, and  $(AB)^* = B^* A^*$  for pseudo-differential operators A and B. The adjoint wave function satisfies the equations

$$L_{mKP}^* \Psi^* = \lambda \Psi^*,$$

$$\partial_{t_k} \Psi^* = -Q_k^* \Psi^*.$$

The Orlov-Schulman operator of the mKP hierarchy is defined by

$$M = \sum_{n=1}^{\infty} n t_n L_{mKP}^{n-1} + \sum_{i=1}^{\infty} v_i L_{mKP}^{-i-1}. \tag{2.2}$$

The constrained mKP hierarchy is

$$\partial_{t_k} L^n = [Q_k, L^n],$$

the Lax operator  $L^n$  has the following relationship,

$$L^n = (L^n)_{\geq 1} + \sum_{i=1}^N v_i(t) \partial^{-1} r_i(t) \partial, \tag{2.3}$$

where  $v_i(t), r_i(t)$  and  $Q_k$  satisfy

$$v_{i,t_k} = Q_k(v_i),$$

$$r_{i,t_k} = -(\partial^{-1} Q_k^* \partial)(r_i), i = 1, 2, 3, \dots, N,$$

$$Q_k = [(L^n)_{\geq 1} + \sum_{i=1}^N v_i(t) \partial^{-1} r_i(t) \partial]^{k/n}_{\geq 1}.$$

The Lax equation of the dispersionless mKP hierarchy is defined by ( $\mathcal{L}_{dmKP} = \lambda$ )

$$\mathcal{L}_{dmKP} = P + V_0 + V_1 P^{-1} + V_2 P^{-2} + \dots, \tag{2.4}$$

$\mathcal{L}_{dmKP}$  satisfies the Lax equations

$$\partial_{T_n} \mathcal{L}_{dmKP} = \{Q_n, \mathcal{L}_{dmKP}\}, \tag{2.5}$$

where  $Q_n(P)$  is defined by

$$Q_n = (\mathcal{L}_{dmKP}^n)_{\geq 1}.$$

The Poisson bracket  $\{, \}$  is defined by

$$\{f(X, P), g(X, P)\} = \frac{\partial f}{\partial P} \frac{\partial g}{\partial X} - \frac{\partial f}{\partial X} \frac{\partial g}{\partial P}. \tag{2.6}$$

The dmKP hierarchy can be also written in the zero-curvature form

$$\frac{\partial Q_n(P)}{\partial T_m} - \frac{\partial Q_m(P)}{\partial T_n} + \{Q_n(P), Q_m(P)\} = 0.$$

We consider the quasiclassical limit of the constrained mKP hierarchy to obtain the constrained dispersionless mKP hierarchy, by taking  $t_n \rightarrow \epsilon t_n = T_n$  [24],

$$L_\epsilon^n = Q_{\epsilon, n} + \sum_{i=1}^N v_i \left(\frac{T}{\epsilon}\right) (\epsilon \partial)^{-1} r_i \left(\frac{T}{\epsilon}\right) \epsilon \partial,$$

$$Q_{\epsilon, n} = (L_\epsilon^n)_{\geq 1},$$

where  $v_i(\frac{T}{\epsilon}), r_i(\frac{T}{\epsilon})$  and  $Q_{\epsilon, k}$  satisfy

$$\epsilon (v_i(\frac{T}{\epsilon}))_{T_k} = Q_{\epsilon, k} (v_i(\frac{T}{\epsilon})),$$

$$\epsilon (r_i(\frac{T}{\epsilon}))_{T_k} = -((\epsilon \partial)^{-1} Q_{\epsilon, k}^* (\epsilon \partial)) (r_i(\frac{T}{\epsilon})), i = 1, 2, 3, \dots, N,$$

$$Q_{\epsilon, k} = ((L_\epsilon^n)^{k/n})_{\geq 1}.$$

It can be proved that  $v_i(\frac{T}{\epsilon})$  and  $r_i(\frac{T}{\epsilon})$  have the following WKB asymptotic expansion as  $\epsilon \rightarrow 0$ ,

$$v_i \left(\frac{T}{\epsilon}\right) \sim \exp\left[\frac{S(T, \lambda_i)}{\epsilon} + a_{i1} + O(\epsilon)\right],$$

$$r_i \left(\frac{T}{\epsilon}\right) \sim \exp\left[-\frac{S(T, \lambda_i)}{\epsilon} + a_{i2} + O(\epsilon)\right].$$

By calculation,

$$v_i \left(\frac{T}{\epsilon}\right) (\epsilon \partial)^{-1} r_i \left(\frac{T}{\epsilon}\right) (\epsilon \partial) = -\frac{e^{a_{i1} + a_{i2}}}{S(T, \lambda_i)_X} \times (1 + (S(T, \lambda_i)_X + O(\epsilon)) (\epsilon \partial)^{-1} + \dots + ((S(T, \lambda_i)_X)^n + O(\epsilon)) (\epsilon \partial)^{-n} + \dots).$$

We can obtain the constrained dispersionless mKP hierarchy for  $\mathcal{L}$  as

$$\mathcal{L}^n = Q_n - \sum_{i=1}^N \left(\frac{a_i}{P_i} + \frac{a_i}{P - P_i}\right), \tag{2.7}$$

$$\mathcal{L} = \{Q_n, \mathcal{L}\}, \tag{2.8}$$

where  $Q_n = (\mathcal{L}^n)_{\geq 1}$ ,  $a_i, P_i$  satisfy

$$a_i = e^{a_{i1} + a_{i2}}, \tag{2.9}$$

$$P_i = S(T, \lambda_i)_X, \tag{2.10}$$

$$\frac{\partial a_i}{\partial T_n} = [a_i \left(\frac{\partial Q_n}{\partial P}\right) |_{P=P_i}]_X, \tag{2.11}$$

$$\frac{\partial P_i}{\partial T_n} = [Q_n |_{P=P_i}]_X. \tag{2.12}$$

When  $n = 1, N = 1$ ,

$$\mathcal{L} = P - \left(\frac{a_1}{P_1} + \frac{a_1}{P - P_1}\right), \tag{2.13}$$

we can get its Taylor expansion

$$\mathcal{L} = P - a_1 \sum_{i=0}^{+\infty} P_1^{i-1} P^{-i}. \tag{2.14}$$

As we all know, the symmetry constraint for Harry Dym hierarchy is given by

$$L_{cDym}^n = B_n + \sum_{i=1}^N q_i \partial^{-1} u_i \partial^2,$$

where  $q_i$ ,  $u_i$  and  $B_n$  satisfy

$$\begin{aligned} \frac{\partial q_i}{\partial t_n} &= B_n(q_i), \\ \frac{\partial u_i}{\partial t_n} &= -\partial^{-2} B_n^* \partial^2(u_i), i = 1, 2, 3, \dots, N, \\ B_n &= (L_{cDym}^n)_{\geq 2}, \end{aligned}$$

where  $B_n^*$  is the adjoint operator of  $B_n$ .

The dispersionless Harry Dym hierarchy is given by

$$\mathcal{L}_{dDym} = U_1 P + U_0 + \frac{U_{-1}}{P} + \frac{U_{-2}}{P^2} + \dots,$$

it satisfies the Lax equations

$$\partial_{T_k} \mathcal{L}_{dDym} = \{\mathcal{B}_k(P), \mathcal{L}_{dDym}\},$$

where

$$\mathcal{B}_k(P) = (\mathcal{L}_{dDym}^k)_{\geq 2}.$$

Using the same method, consider the quasiclassical limit of the constrained Harry Dym hierarchy, we can obtain the constrained dispersionless Harry Dym hierarchy [20],

$$\mathcal{L}^n = \mathcal{B}_n + \sum_{i=1}^N \left( \frac{\alpha_i P}{\beta_i^2} + \frac{\alpha_i}{\beta_i} + \frac{\alpha_i}{P - \beta_i} \right), \tag{2.15}$$

$$\mathcal{L}_{T_n} = \{\mathcal{B}_n, \mathcal{L}\}, \tag{2.16}$$

where  $\mathcal{B}_n = (\mathcal{L}^n)_{\geq 2}$ ,  $\alpha_i, \beta_i$  satisfy

$$\alpha_i = e^{\alpha_{i1} + \alpha_{i2}}, \tag{2.17}$$

$$\beta_i = S(T, \lambda_i)_X, \tag{2.18}$$

$$\frac{\partial \alpha_i}{\partial T_n} = [\alpha_i \left( \frac{\partial \mathcal{B}_n}{\partial P} \right) |_{P=\beta_i}]_X, \tag{2.19}$$

$$\frac{\partial \beta_i}{\partial T_n} = [\mathcal{B}_n |_{P=\beta_i}]_X. \tag{2.20}$$

### 3. Auto-Bäcklund transformations for dmKPH

In section 3, we introduce the auto-Bäcklund transformations for dmKP hierarchy by using the method in [19].

We have

$$\lambda = P^m + a_{m-1} P^{m-1} + a_{m-2} P^{m-2} + \dots + a_1 P + a_0 + a_{-1} P^{-1} + a_{-2} P^{-2} + \dots,$$

where  $a_{m-1}, a_{m-2}, \dots, a_1, a_0, a_{-1}, a_{-2}, \dots$  are functions of  $T = (T_1 = x, T_2, T_3, \dots)$ .  $\phi(T)$  is arbitrary function of  $T$ , and it is independent of  $P$ .

We define

$$\tilde{\lambda} = e^{-ad\phi} \lambda = \lambda - \{\phi, \lambda\} + \frac{1}{2} \{\phi, \{\phi, \lambda\}\} - \frac{1}{3} \{\phi, \{\phi, \{\phi, \lambda\}\}\} + \dots,$$

where the Poisson bracket is defined by (2.6). By calculation, we can obtain

$$\tilde{\lambda} = \sum_{n=0}^{\infty} \frac{1}{n!} (\phi_x)^n \partial_P^n \lambda.$$

**Lemma 3.1.** *The following identities hold true [19]*

$$\begin{aligned}
 e^{-ad\phi}(\mu_1\mu_2) &= e^{-ad\phi}\mu_1 \cdot e^{-ad\phi}\mu_2, \\
 \tilde{\mu}_{\geq 1} &= e^{-ad\phi}(\mu_{\geq 0}) - \mu_{\geq 0}|_{P=\phi_x} \\
 &= \mu_{\geq 0} + \phi_x\partial_P(\mu_{\geq 0}) + \frac{1}{2!}\phi_x^2\partial_P^2(\mu_{\geq 0}) + \dots \\
 &\quad - (\phi_x^m + a_{m-1}\phi_x^{m-1} + \dots + a_1\phi_x + a_0) \\
 &= e^{-ad\phi}(\mu_{\geq 1}) - \mu_{\geq 1}|_{P=\phi_x}, \\
 \tilde{\mu}_{T_q} - \{(\tilde{\mu}^n)_{\geq 1}, \tilde{\mu}\} &= e^{-ad\phi}(\mu_{T_q} - \{(\mu^n)_{\geq 0}, \mu\}) - \{\phi_{T_q} - (\mu^n)_{\geq 0}|_{P=\phi_x}, \tilde{\mu}\} \\
 &= e^{-ad\phi}(\mu_{T_q} - \{(\mu^n)_{\geq 1}, \mu\}) - \{\phi_{T_q} - (\mu^n)_{\geq 1}|_{P=\phi_x}, \tilde{\mu}\}.
 \end{aligned}$$

Let

$$\mu = P + V_0 + V_1P^{-1} + V_2P^{-2} + \dots,$$

we suppose that  $V_i(T)$  satisfy the dmKP hierarchy in (2.5), and  $\phi(T)$  satisfies the following equation,

$$\phi_{T_q} = (\mu^n)_{\geq 1}|_{P=\phi_x}.$$

Then, we have  $\tilde{\mu} = e^{-ad\phi}\mu$  also satisfy the dmKP hierarchy.

$$\tilde{\mu} = P + \tilde{V}_0 + \tilde{V}_1P^{-1} + \tilde{V}_2P^{-2} + \dots,$$

we can get  $\tilde{V}_i = V_i$ , it is the auto-Bäcklund transformations for the dispersionless mKP hierarchy.

#### 4. Dispersionless Miura map between the constrained dispersionless mKP hierarchy and the constrained dispersionless Harry Dym hierarchy

In section 4, we introduce the dispersionless Miura map between the constrained dispersionless mKP hierarchy and the constrained dispersionless Harry Dym hierarchy by using the method in [20].

**Lemma 4.1.** *Let [20]*

$$x' = \psi, T'_n = T_n, \mathcal{L}(x', T') = \mathcal{L}(x, T), P' = \psi_x^{-1}P.$$

Noting that,

$$\begin{aligned}
 \frac{\partial}{\partial T_n} &= \frac{\partial T'_n}{\partial T_n} \frac{\partial}{\partial T'_n} + \frac{\partial x'}{\partial T_n} \frac{\partial}{\partial x'} = \frac{\partial}{\partial T'_n} + \frac{\partial \psi}{\partial T_n} \frac{\partial}{\partial x'}, \\
 A'_{\geq 2} &= A'_{\geq 1} - A'_{[1]} = A'_{\geq 1} - \{A'_{\geq 1}, x'\}_{[0]}P' = A_{\geq 1} - \{A_{\geq 1}, \psi\}_{[0]}P',
 \end{aligned}$$

we have

$$\begin{aligned}
 \mathcal{L}_{T'_n} &= \frac{\partial \mathcal{L}}{\partial T_n} - \frac{\partial \psi}{\partial T_n} \frac{\partial \mathcal{L}}{\partial x'} = \mathcal{L}_{T_n} - \left\{ \frac{\partial \psi}{\partial T_n} P', \mathcal{L} \right\}, \\
 \mathcal{B}_n &= \mathcal{Q}_n - \{ \mathcal{Q}_n, \psi \}_{[0]}P'.
 \end{aligned}$$

We can obtain

$$\begin{aligned}
 \mathcal{L}_{T_n} - \{ \mathcal{B}_n, \mathcal{L} \} &= \mathcal{L}_{T_n} - \left\{ \frac{\partial \psi}{\partial T_n} P', \mathcal{L} \right\} - \{ \mathcal{Q}_n - \{ \mathcal{Q}_n, \psi \}_{[0]}P', \mathcal{L} \} \\
 &= \mathcal{L}_{T_n} - \{ \mathcal{Q}_n, \mathcal{L} \} - \left\{ \left( \frac{\partial \psi}{\partial T_n} - \{ \mathcal{Q}_n, \psi \}_{[0]} \right) P', \mathcal{L} \right\}.
 \end{aligned}$$

**Theorem 4.1.** *If  $\mathcal{L}, a_i, P_i$  satisfy the constrained dispersionless mKP hierarchy (2.14) and  $\psi$  is a function of  $(x, T)$  satisfying*

$$\frac{\partial \psi}{\partial T_n} = \{ \mathcal{Q}_n, \psi \}_{[0]},$$

then  $\mathcal{L}$  satisfy the constrained dispersionless Harry Dym hierarchy (2.15).

**Proof.** The Lemma (4.1) indicates that  $\mathcal{L}(x, T) = \mathcal{L}(x', T')$ . Then, using the same method as in [20], we can indicate that  $\alpha_i, \beta_i$  satisfy the constrained dispersionless Harry Dym hierarchy through these relationships

$$\begin{aligned}
 \alpha_i &= -\psi_x^{-1}a_i, \\
 \beta_i &= \psi_x^{-1}P_i,
 \end{aligned}$$

which have been confirmed in [22].  $\square$

### 5. The additional symmetries of the constrained dispersionless mKP hierarchy

In section 5, we introduce the additional symmetries of the constrained dispersionless mKP hierarchy by introducing a new Laurent series  $\mathcal{Y}$ .

The additional symmetries of the dispersionless mKP hierarchy were given by

$$\frac{\partial \mathcal{L}_{dmKP}}{\partial t_{l,m}^*} = -\{(\mathcal{M}^m \mathcal{L}_{dmKP}^l)_{<1}, \mathcal{L}_{dmKP}\},$$

or they can be written as

$$\frac{\partial \mathcal{L}_{dmKP}}{\partial t_{l,m}^*} = \{(\mathcal{M}^m \mathcal{L}_{dmKP}^l)_{\geq 1}, \mathcal{L}_{dmKP}\} + m \mathcal{M}^{m-1} \mathcal{L}_{dmKP}^l. \tag{5.1}$$

The function  $\mathcal{M}$  satisfies the equations

$$\frac{\partial \mathcal{M}}{\partial t_{l,m}^*} = -\{(\mathcal{M}^m \mathcal{L}_{dmKP}^l)_{<1}, \mathcal{M}\},$$

where the Orlov-Schulman function of the dispersionless mKP hierarchy is defined by [31]

$$\mathcal{M} = \sum_{n=1}^{+\infty} n T_n \mathcal{L}_{dmKP}^{n-1} + \sum_{i=1}^{+\infty} V_i \mathcal{L}_{dmKP}^{-i-1}, \tag{5.2}$$

we have  $\{\mathcal{M}, \mathcal{L}_{dmKP}\} = -1$ .

Furthermore, we have

$$\frac{\partial \mathcal{M}^n \mathcal{L}_{dmKP}^k}{\partial t_{l,m}^*} = -\{(\mathcal{M}^m \mathcal{L}_{dmKP}^l)_{<1}, \mathcal{M}^n \mathcal{L}_{dmKP}^k\}.$$

Now, we apply the additional symmetric flows defined by (5.1) to  $\mathcal{L}$  which is defined by (2.14). When  $m = 1$ , we have

$$\left(\frac{\partial \mathcal{L}}{\partial t_{l,1}^*}\right)_{<1} = \{(\mathcal{M} \mathcal{L}^l)_{\geq 1}, \mathcal{L}\}_{<1} + (\mathcal{L}^l)_{<1}. \tag{5.3}$$

Next let us analyze the two terms  $\{(\mathcal{M} \mathcal{L}^l)_{\geq 1}, \mathcal{L}\}_{<1}, (\mathcal{L}^l)_{<1}$ .

**Theorem 5.1.** *The Laurent series  $\mathcal{L}$  and the Orlov function  $\mathcal{M}$  of the constrained dispersionless mKP hierarchy have the following relationship.*

$$\begin{aligned} ((\mathcal{M} \mathcal{L}^l)_{\geq 1}, \mathcal{L})_{<1} &= -\frac{\partial a_1}{\partial x} \sum_{i=0}^{+\infty} \sum_{n=1}^{+\infty} n (\mathcal{M} \mathcal{L}^l)_{[n]} P_1^{i+n-2} P^{-i} \\ &\quad - a_1 \frac{\partial P_1}{\partial x} \sum_{i=0}^{+\infty} \sum_{n=1}^{+\infty} n(n+i-2) (\mathcal{M} \mathcal{L}^l)_{[n]} P_1^{n+i-3} P^{-i} \\ &\quad - a_1 \sum_{i=0}^{+\infty} \sum_{n=1}^{+\infty} (n+i-1) \frac{\partial (\mathcal{M} \mathcal{L}^l)_{[n]}}{\partial x} P_1^{n+i-2} P^{-i}, \end{aligned} \tag{5.4}$$

we define that  $(A)_{[i]}$  mean  $A_i$  for arbitrary formal Laurent series  $A = \sum_i A_i P^{-i}$ .

**Proof.** By using the equation (2.6) and (2.14)

$$\begin{aligned} \{(\mathcal{M} \mathcal{L}^l)_{\geq 1}, \mathcal{L}\}_{<1} &= \frac{\partial (\mathcal{M} \mathcal{L}^l)_{\geq 1}}{\partial P} \frac{\partial \mathcal{L}_{<1}}{\partial X} - \frac{\partial (\mathcal{M} \mathcal{L}^l)_{\geq 1}}{\partial X} \frac{\partial \mathcal{L}_{<1}}{\partial P} \\ &= \left(\frac{\partial (\mathcal{M} \mathcal{L}^l)_{\geq 1}}{\partial P} \left(-\frac{\partial a_1}{\partial X} \sum_{i=0}^{+\infty} P_1^{i-1} P^{-i} - a_1 \frac{\partial P_1}{\partial X} \sum_{i=0}^{+\infty} (i-1) P_1^{i-2} P^{-i}\right)\right)_{<1} \\ &\quad - \left(\frac{\partial (\mathcal{M} \mathcal{L}^l)_{\geq 1}}{\partial X} a_1 \sum_{i=0}^{+\infty} i P_1^{i-1} P^{-i-1}\right)_{<1} \\ &= -\sum_{n=1}^{+\infty} \sum_{i=n-1}^{+\infty} n (\mathcal{M} \mathcal{L}^l)_{[n]} \frac{\partial a_1}{\partial X} P_1^{i-1} P^{-i+n-1} - \sum_{n=1}^{+\infty} \sum_{i=n-1}^{+\infty} (i-1) n (\mathcal{M} \mathcal{L}^l)_{[n]} a_1 \frac{\partial P_1}{\partial X} P_1^{i-2} P^{-i+n-1} \\ &\quad - a_1 \sum_{n=1}^{+\infty} \sum_{i=n-1}^{+\infty} \frac{\partial (\mathcal{M} \mathcal{L}^l)_{[n]}}{\partial X} P_1^{i-1} i P^{-i+n-1} \end{aligned}$$

$$\begin{aligned}
 &= -\frac{\partial a_1}{\partial x} \sum_{i=0}^{+\infty} \sum_{n=1}^{+\infty} n(\mathcal{ML}^l)_{[n]} P_1^{i+n-2} P^{-i} - a_1 \frac{\partial P_1}{\partial x} \sum_{i=0}^{+\infty} \sum_{n=1}^{+\infty} n(n+i-2)(\mathcal{ML}^l)_{[n]} P_1^{n+i-3} P^{-i} \\
 &\quad - a_1 \sum_{i=0}^{+\infty} \sum_{n=1}^{+\infty} (n+i-1) \frac{\partial(\mathcal{ML}^l)_{[n]}}{\partial x} P_1^{n+i-2} P^{-i}. \quad \square
 \end{aligned}$$

**Theorem 5.2.** *The Laurent series  $\mathcal{L}$  of the constrained dispersionless mKP hierarchy has the following relationship.*

$$(\mathcal{L}^l)_{<1} = \sum_{k=1}^l C_l^k \mathcal{L}_{l-k} \mathcal{L}_0^{k-1}, \quad l = 1, 2, 3, \dots, \tag{5.5}$$

$$(\mathcal{L}^0)_{<1} = 1. \tag{5.6}$$

We define that

$$\mathcal{L}_j = -a_1 \sum_{i=j}^{+\infty} P_1^{i-1} P^{-i+j}.$$

For  $l = 0, 1, 2$ , the flows can be rewritten as

$$\left(\frac{\partial \mathcal{L}}{\partial t_{l,1}^*}\right)_{<1} = -\frac{\partial a_1}{\partial t_{l,1}^*} \sum_{i=0}^{+\infty} P_1^{i-1} P^{-i} - a_1 \frac{\partial P_1}{\partial t_{l,1}^*} \sum_{i=0}^{+\infty} (i-1) P_1^{i-2} P^{-i}.$$

When  $l = 0$ ,

$$\left(\frac{\partial \mathcal{L}}{\partial t_{0,1}^*}\right)_{<1} = -\frac{\partial a_1}{\partial t_{0,1}^*} \sum_{i=0}^{+\infty} P_1^{i-1} P^{-i} - a_1 \frac{\partial P_1}{\partial t_{0,1}^*} \sum_{i=0}^{+\infty} (i-1) P_1^{i-2} P^{-i},$$

using the equation (5.3),

$$\begin{aligned}
 \left(\frac{\partial \mathcal{L}}{\partial t_{0,1}^*}\right)_{<1} &= -\frac{\partial a_1}{\partial x} \sum_{i=0}^{+\infty} \sum_{n=1}^{+\infty} n(\mathcal{ML}^0)_{[n]} P_1^{i+n-2} P^{-i} \\
 &\quad - a_1 \frac{\partial P_1}{\partial x} \sum_{i=0}^{+\infty} \sum_{n=1}^{+\infty} n(n+i-2)(\mathcal{ML}^0)_{[n]} P_1^{n+i-3} P^{-i} \\
 &\quad - a_1 \sum_{i=0}^{+\infty} \sum_{n=1}^{+\infty} (n+i-1) \frac{\partial(\mathcal{ML}^0)_{[n]}}{\partial x} P_1^{n+i-2} P^{-i} + 1.
 \end{aligned}$$

When  $l = 1$ ,

$$\left(\frac{\partial \mathcal{L}}{\partial t_{1,1}^*}\right)_{<1} = -\frac{\partial a_1}{\partial t_{1,1}^*} \sum_{i=0}^{\infty} P_1^{i-1} P^{-i} - a_1 \frac{\partial P_1}{\partial t_{1,1}^*} \sum_{i=0}^{\infty} (i-1) P_1^{i-2} P^{-i},$$

using the equation (5.3),

$$\begin{aligned}
 \left(\frac{\partial \mathcal{L}}{\partial t_{1,1}^*}\right)_{<1} &= -\frac{\partial a_1}{\partial x} \sum_{i=0}^{+\infty} \sum_{n=1}^{+\infty} n(\mathcal{ML}^1)_{[n]} P_1^{i+n-2} P^{-i} \\
 &\quad - a_1 \frac{\partial P_1}{\partial x} \sum_{i=0}^{+\infty} \sum_{n=1}^{+\infty} n(n+i-2)(\mathcal{ML}^1)_{[n]} P_1^{n+i-3} P^{-i} \\
 &\quad - a_1 \sum_{i=0}^{+\infty} \sum_{n=1}^{+\infty} (n+i-1) \frac{\partial(\mathcal{ML}^1)_{[n]}}{\partial x} P_1^{n+i-2} P^{-i} + \mathcal{L}_0.
 \end{aligned}$$

When  $l = 2$ ,

$$\left(\frac{\partial \mathcal{L}}{\partial t_{2,1}^*}\right)_{<1} = -\frac{\partial a_1}{\partial t_{2,1}^*} \sum_{i=0}^{\infty} P_1^{i-1} P^{-i} - a_1 \frac{\partial P_1}{\partial t_{2,1}^*} \sum_{i=0}^{\infty} (i-1) P_1^{i-2} P^{-i},$$

using the equation (5.3),

$$\begin{aligned} \left(\frac{\partial \mathcal{L}}{\partial t_{2,1}^*}\right)_{<1} &= -\frac{\partial a_1}{\partial x} \sum_{i=0}^{+\infty} \sum_{n=1}^{+\infty} n(\mathcal{ML}^2)_{[n]} P_1^{i+n-2} P^{-i} \\ &\quad - a_1 \frac{\partial P_1}{\partial x} \sum_{i=0}^{+\infty} \sum_{n=1}^{+\infty} n(n+i-2)(\mathcal{ML}^2)_{[n]} P_1^{n+i-3} P^{-i} \\ &\quad - a_1 \sum_{i=0}^{+\infty} \sum_{n=1}^{+\infty} (n+i-1) \frac{\partial(\mathcal{ML}^2)_{[n]}}{\partial x} P_1^{n+i-2} P^{-i} + \mathcal{L}_0^2 + 2\mathcal{L}_1. \end{aligned}$$

We can find that the flows do not hold for  $l = 0, 1, 2, \dots$

Therefore, we construct new flows,

$$\left(\frac{\partial \mathcal{L}}{\partial t_{l,1}^*}\right)_{<1} = \{-(\mathcal{ML}^l)_{<1} + \mathcal{Y}_l, \mathcal{L}\}_{<1}, \tag{5.7}$$

where

$$\mathcal{Y}_l = -a_1 \sum_{k=1}^l C_l^k \sum_{i=l-k}^{+\infty} \frac{P_1^{i-1} P^{-i+l-k+1}}{-i+l-k+1}, l = 1, 2, 3, \dots, \tag{5.8}$$

$$\mathcal{Y}_0 = \mathcal{M}, \tag{5.9}$$

in which  $a_1, P_1$  satisfy the following equations

$$-\frac{\partial a_1}{\partial x} \sum_{i=0}^{+\infty} P_1^{i-1} P^{-i} - a_1 \frac{\partial P_1}{\partial x} \sum_{i=0}^{+\infty} (i-1) P_1^{i-2} P^{-i} + \sum_{k_1=1}^l \mathcal{L}_0^{k_1-1} = 0, k_1 = k.$$

We define that

$$\begin{aligned} \sum_{k=1}^l \sum_{i=0}^{+\infty} i^{-1} \left(\frac{\partial a_1}{\partial t_j^*} P_1^i + a_1 \frac{\partial P_1}{\partial t_j^*} (l-k+i) P_1^{i-1}\right) &= -\sum_{k=1}^l \sum_{n=1}^{+\infty} \sum_{i=0}^{+\infty} \left(\frac{\partial a_1}{\partial x} \frac{n(\mathcal{ML}^j)_{[n]} P_1^{n+i-1}}{-n-i+1}\right. \\ &\quad \left.+ a_1 \frac{\partial P_1}{\partial x} \frac{(l-k+n+i-1)n(\mathcal{ML}^j)_{[n]} P_1^{n+i-2}}{-n-i+1}\right. \\ &\quad \left.+ a_1 P_1^{n+i-1} \frac{\partial(\mathcal{ML}^j)_{[n]}}{\partial x}\right) + (l-j)a_1 \sum_{i=0}^{+\infty} i^{-1} P_1^{j+i-1}. \end{aligned}$$

**Lemma 5.1.** The Laurent series  $\mathcal{L}$  and  $\mathcal{Y}_l$  satisfy the following relation,

$$\begin{aligned} \{\mathcal{Y}_l, \mathcal{L}\}_{<1} &= -(\mathcal{L}^l)_{<1} + a_1 \sum_{k=1}^l \sum_{m=0}^{+\infty} \sum_{i=0}^{+\infty} \frac{i C_l^k}{-i+1} \left(\frac{\partial a_1}{\partial x} P_1^{l-k+i+m-2}\right. \\ &\quad \left.+ a_1 \frac{\partial P_1}{\partial x} (l-k+m-1) P_1^{l-k+i+m-3}\right) P^{-i-m}, l = 1, 2, 3, \dots, \\ \{\mathcal{Y}_0, \mathcal{L}\}_{<1} &= -1. \end{aligned}$$

The Laurent series  $\mathcal{Y}_l$  also have the following relationship.

**Lemma 5.2.** The formal Laurent series  $\mathcal{Y}_l$  satisfy the following relation,

$$\frac{\partial \mathcal{Y}_l}{\partial t_j^*} = \{(\mathcal{ML}^j)_{\geq 1}, \mathcal{Y}_l\}_{<1} + (l-j)\mathcal{Y}_{l+j-1}, \tag{5.10}$$

where  $j, l = 0, 1, 2, \dots$

**Proof.** The left-hand side of equation (5.10) is

$$\frac{\partial \mathcal{Y}_l}{\partial t_j^*} = -\sum_{k=1}^l \sum_{i=l-k}^{+\infty} \frac{C_l^k}{-i+l-k+1} \left(\frac{\partial a_1}{\partial t_j^*} P_1^{i-1} + a_1 \frac{\partial P_1}{\partial t_j^*} (i-1) P_1^{i-2}\right) P^{-i+l-k+1},$$

the first term on the right-hand side of equation (5.10) is

$$\begin{aligned} \{(\mathcal{ML}^j)_{\geq 1}, \mathcal{Y}_l\}_{<1} = & -\sum_{n=1}^{+\infty} \sum_{k=1}^l \sum_{i=l-k+n}^{+\infty} C_l^k \left( \frac{\partial a_1}{\partial x} \frac{n(\mathcal{ML}^j)_{[n]} P_1^{i-1}}{-i+l-k+1} + a_1 \frac{\partial P_1}{\partial x} \frac{(i-1)n(\mathcal{ML}^j)_{[n]} P_1^{i-2}}{-i+l-k+1} \right. \\ & \left. + a_1 P_1^{i-1} \frac{\partial(\mathcal{ML}^j)_{[n]}}{\partial x} \right) P^{-i+l-k+n}, \end{aligned}$$

the second term on the right-hand side of equation (5.10) is

$$(l-j)\mathcal{Y}_{l+j-1} = -(l-j)a_1 \sum_{k=1}^l \sum_{i=l+j-k-1}^{+\infty} \frac{C_l^k}{-i+l+j-k} P_1^{i-1} P^{-i+l-j-k}.$$

By comparison, we know that the left and right sides of the equation (5.10) are equal.  $\square$

**Lemma 5.3.** *The additional flows  $\frac{\partial}{\partial t_l^*}$  commute with the flows  $\frac{\partial}{\partial t_n}$  of the constrained dispersionless mKP hierarchy.*

**Proof.**

$$\begin{aligned} \left[ \frac{\partial}{\partial t_l^*}, \frac{\partial}{\partial t_n} \right] \mathcal{L} &= \frac{\partial}{\partial t_l^*} \{ \mathcal{B}_n, \mathcal{L} \} - \frac{\partial}{\partial t_n} \{ -(\mathcal{ML}^l)_{<1} + \mathcal{Y}_l, \mathcal{L} \} \\ &= \left\{ \frac{\partial \mathcal{B}_n}{\partial t_l^*}, \mathcal{L} \right\} + \left\{ \mathcal{B}_n, \frac{\partial \mathcal{L}}{\partial t_l^*} \right\} - \left\{ \frac{\partial(-\mathcal{ML}^l + \mathcal{Y}_l)_{<1}}{\partial t_n}, \mathcal{L} \right\} - \{ (-\mathcal{ML}^l + \mathcal{Y}_l)_{<1}, \frac{\partial \mathcal{L}}{\partial t_n} \} \\ &= \{ \{-\mathcal{ML}^l + \mathcal{Y}_l\}_{<1}, (\mathcal{L}^n)_{\geq 1} \}_{\geq 1}, \mathcal{L} \} + \{ (\mathcal{L}^n)_{\geq 1}, \{ (-\mathcal{ML}^l + \mathcal{Y}_l)_{<1}, \mathcal{L} \} \} \\ &\quad - \{ \{ (\mathcal{L}^n)_{\geq 1}, (-\mathcal{ML}^l + \mathcal{Y}_l)_{<1} \}_{<1}, \mathcal{L} \} - \{ (-\mathcal{ML}^l + \mathcal{Y}_l)_{<1}, \{ (\mathcal{L}^n)_{\geq 1}, \mathcal{L} \} \} \\ &= \{ \{-\mathcal{ML}^l + \mathcal{Y}_l\}_{<1}, (\mathcal{L}^n)_{\geq 1} \}, \mathcal{L} \} + \{ (\mathcal{L}^n)_{\geq 1}, \{ (-\mathcal{ML}^l + \mathcal{Y}_l)_{<1}, \mathcal{L} \} \} \\ &\quad - \{ (-\mathcal{ML}^l + \mathcal{Y}_l)_{<1}, \{ (\mathcal{L}^n)_{\geq 1}, \mathcal{L} \} \}. \end{aligned}$$

By using the Jacobi identity of the Poisson bracket  $\{ \{ f, g \}, h \} + \{ \{ g, h \}, f \} + \{ \{ h, f \}, g \} = 0$ , we can get the last equation is 0.  $\square$

**Lemma 5.4.** *The additional flows act on the space of the formal Laurent series  $\mathcal{L}$  of the constrained dispersionless mKP hierarchy, forming a subalgebra of the Virasoro algebra.*

**Proof.**

$$\begin{aligned} \left[ \frac{\partial}{\partial t_l^*}, \frac{\partial}{\partial t_j^*} \right] \mathcal{L} &= \frac{\partial}{\partial t_l^*} \{ (-\mathcal{ML}^j)_{<1}, \mathcal{L} \} - \frac{\partial}{\partial t_j^*} \{ (-\mathcal{ML}^l + \mathcal{Y}_l)_{<1}, \mathcal{L} \} \\ &= \left\{ \frac{\partial(-\mathcal{ML}^j)_{<1}}{\partial t_l^*}, \mathcal{L} \right\} + \{ (-\mathcal{ML}^j)_{<1}, \frac{\partial \mathcal{L}}{\partial t_l^*} \} \\ &\quad - \left\{ \frac{\partial(-\mathcal{ML}^l + \mathcal{Y}_l)_{<1}}{\partial t_j^*}, \mathcal{L} \right\} - \{ (-\mathcal{ML}^l + \mathcal{Y}_l)_{<1}, \frac{\partial \mathcal{L}}{\partial t_j^*} \} \\ &= \{ \{ (-\mathcal{ML}^l + \mathcal{Y}_l)_{<1}, (-\mathcal{ML}^j)_{<1} \}_{<1}, \mathcal{L} \} \\ &\quad - \{ \{ (-\mathcal{ML}^j)_{<1}, (-\mathcal{ML}^l + \mathcal{Y}_l)_{<1} \}_{<1}, \mathcal{L} \} \\ &\quad - \{ \{ (-\mathcal{ML}^l + \mathcal{Y}_l)_{<1}, (-\mathcal{ML}^j)_{<1} \}_{<1}, \mathcal{L} \} \\ &= \{ \{-\mathcal{ML}^l + \mathcal{Y}_l, -\mathcal{ML}^j\}_{<1}, \mathcal{L} \}. \end{aligned}$$

By the equation (5.10), we can get

$$\{ -\mathcal{ML}^l + \mathcal{Y}_l, -\mathcal{ML}^j \}_{<1} = (j-l) \{ -(\mathcal{ML}^{j+l-1})_{<1} + \mathcal{Y}_{j+l-1} \},$$

it also means that

$$\left[ \frac{\partial}{\partial t_l^*}, \frac{\partial}{\partial t_j^*} \right] = (j-l) \frac{\partial}{\partial t_{j+l-1}^*},$$

where  $j, l \in \mathbb{Z}_{\geq 0}$ .  $\square$

**Theorem 5.3.** *The additional flows acting on  $a_1$  and  $P_1$  of the constrained dispersionless mKP hierarchy are given by*

$$\begin{aligned} \frac{\partial a_1}{\partial t_{l,1}^*} &= \frac{\partial a_1}{\partial x} \sum_{n=1}^{+\infty} n(\mathcal{ML})_{[n]} P_1^{n-1}, \\ \frac{\partial P_1}{\partial t_{l,1}^*} &= -\left( \frac{\partial P_1}{\partial x} \sum_{n=1}^{+\infty} n(n-2)(\mathcal{ML})_{[n]} P_1^{n-1} + \sum_{n=1}^{+\infty} (n-1) \frac{\partial(\mathcal{ML})_{[n]}}{\partial x} P_1^n \right). \end{aligned}$$

When  $l = 0$ ,

$$\frac{\partial a_1}{\partial t_{0,1}^*} = \frac{\partial a_1}{\partial x} \sum_{n=1}^{+\infty} n \mathcal{M}_{[n]} P_1^{n-1},$$

$$\frac{\partial P_1}{\partial t_{0,1}^*} = -\left(\frac{\partial P_1}{\partial x} \sum_{n=1}^{+\infty} n(n-2) \mathcal{M}_{[n]} P_1^{n-1} + \sum_{n=1}^{+\infty} (n-1) \frac{\partial \mathcal{M}_{[n]}}{\partial x} P_1^n\right).$$

When  $l = 1$ ,

$$\frac{\partial a_1}{\partial t_{1,1}^*} = \frac{\partial a_1}{\partial x} \sum_{n=1}^{+\infty} n (\mathcal{ML})_{[n]} P_1^{n-1},$$

$$\frac{\partial P_1}{\partial t_{1,1}^*} = -\left(\frac{\partial P_1}{\partial x} \sum_{n=1}^{+\infty} n(n-2) (\mathcal{ML})_{[n]} P_1^{n-1} + \sum_{n=1}^{+\infty} (n-1) \frac{\partial (\mathcal{ML})_{[n]}}{\partial x} P_1^n\right).$$

These can be proven by the above lemmas and the equation (5.3).

In (2.7), when  $n = 1$ ,

$$\mathcal{L} = P - \sum_{j=1}^N \left(\frac{a_j}{P_j} + \frac{a_j}{P - P_j}\right),$$

we can get its Taylor expansion

$$\mathcal{L} = P - \sum_{j=1}^N a_j \sum_{i=0}^{+\infty} P_j^{i-1} P^{-i}.$$

Now, we can construct the additional symmetries of the constrained dispersionless mKP hierarchy,

$$\left(\frac{\partial \mathcal{L}}{\partial t_{l,1}^*}\right)_{<1} = \{-(\mathcal{ML}^l)_{<1} + \mathcal{Y}_l^N, \mathcal{L}\}_{<1},$$

where

$$\mathcal{Y}_0^N = \mathcal{M},$$

$$\mathcal{Y}_l^N = \sum_{j=1}^N \left(-a_j \sum_{k=1}^l C_l^k \sum_{i=l-k}^{+\infty} \frac{P_j^{i-1} P^{-i+l-k+1}}{-i+l-k+1}\right), l = 1, 2, 3, \dots,$$

in which  $a_j, P_j$  satisfy the following equations

$$-\sum_{j=1}^N \frac{\partial a_j}{\partial x} \sum_{i=0}^{+\infty} P_j^{i-1} P^{-i} - \sum_{j=1}^N a_j \frac{\partial P_j}{\partial x} \sum_{i=0}^{+\infty} (i-1) P_j^{i-2} P^{-i} + \sum_{k_1=1}^l \mathcal{L}_{j,0}^{k_1-1} = 0, k_1 = k.$$

We define that

$$\mathcal{L}_{j,k} = \sum_{j=1}^N \left(-a_j \sum_{i=k}^{+\infty} P_j^{i-1} P^{-i+k}\right).$$

The Laurent series  $\mathcal{L}$  of the constrained dispersionless mKP hierarchy has the following relationship,

$$(\mathcal{L}^l)_{<1} = \sum_{k=1}^l C_l^k \mathcal{L}_{j,l-k} \mathcal{L}_{j,0}^{k-1}, l = 1, 2, 3, \dots,$$

$$(\mathcal{L}^0)_{<1} = 1.$$

The additional symmetries of the constrained dispersionless mKP hierarchy are constructed.

### 6. Conclusions and discussions

In this article, we have discussed the auto-Bäcklund transformations for dmKP hierarchy, its construction method draws inspiration from the article [19]. In section 4, we construct the dispersionless Miura map between the constrained dispersionless mKP hierarchy and the constrained dispersionless Harry Dym hierarchy. In section 5, we construct the additional symmetries of the constrained dispersionless mKP hierarchy by introducing a new Laurent series  $\mathcal{Y}_l$  in equation (5.8) and (5.9). We have been indicated that the additional flows commute with the flows of the constrained dispersionless mKP hierarchy. We also have been indicated that the additional flows act on the space of the formal Laurent series  $\mathcal{L}$  of the constrained dispersionless mKP hierarchy, forming a subalgebra of the Virasoro algebra. The additional flows acting on  $a_1$  and  $P_1$  of the constrained dispersionless mKP hierarchy are given.

## Declaration of competing interest

We declare that we have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

## Data availability

No data was used for the research described in the article.

## Acknowledgements

This work is supported by the National Natural Science Foundation of China under Grants No. 12071237.

## References

- [1] E. Date, M. Kashiwara, M. Jimbo, T. Miwa, Transformation groups for soliton equations, in: *Nonlinear Integrable Systems—Classical Theory and Quantum Theory*, Kyoto, 1981, World Sci., 1983, pp. 39–119.
- [2] L.A. Dickey, On additional symmetries of the KP hierarchy and Sato's Backlund transformation, *Commun. Math. Phys.* 167 (1995) 227–233.
- [3] C.Z. Li, B. Shou, Supersymmetric quantum spin chains and modified universal characters, *J. Stat. Phys.* 190 (2023) 55.
- [4] Q. Yang, C.Z. Li, Multiparameter universal characters of B-type and integrable hierarchy, *J. Math. Phys.* 63 (2022) 113503.
- [5] C.Z. Li, Multi-component universal character hierarchy and its polynomial tau-functions, *Physica D* 432 (2022) 133166.
- [6] C.Z. Li,  $N = 2$  supersymmetric BKP hierarchy with  $SW_{1+\infty}$  symmetries and its multicomponent generalization, *Phys. Lett. B* 820 (2021) 136563.
- [7] C.Z. Li, Multi-component supersymmetric D type Drinfeld-Sokolov hierarchy and its Virasoro symmetry, *Phys. Lett. B* 823 (2021) 136774.
- [8] C.Z. Li,  $SW_{1+\infty}$  symmetries of  $N = 2$  supersymmetric CKP hierarchy and its multicomponent generalization, *Nucl. Phys. B* 969 (2021) 115465.
- [9] Y. Cheng, M.H. Li, K.L. Tian, On the modified KP hierarchy: tau functions, squared eigenfunction symmetries and additional symmetries, *J. Geom. Phys.* 134 (2018) 19–37.
- [10] K. Takasaki, T. Takebe, Integrable hierarchies and dispersionless limit, *Rev. Math. Phys.* 07 (1995) 743–808.
- [11] J.C. Shaw, M.H. Tu, Miura and auto-Bäcklund transformations for the cKP and cmKP hierarchies, *J. Math. Phys.* 27 (1997) 5756.
- [12] J.P. Cheng, J.S. He, The integral type gauge transformation and the additional symmetry for the constrained KP hierarchy, *Acta Math. Sci.* 35 (2015) 1111–1121.
- [13] H.Z. Chen, L.M. Geng, N. Li, J.P. Cheng, Solving the constrained modified KP hierarchy by gauge transformations, *J. Nonlinear Math. Phys.* 26 (2019) 54–68.
- [14] J.P. Cheng, The gauge transformation of the modified KP hierarchy, *J. Nonlinear Math. Phys.* 25 (2018) 66–85.
- [15] R. Ilangovane, K. Krishnakumar, K.M. Tamizhmani, Algebraic structures on the flows of dispersionless modified KP equation, *Pramana J. Phys.* 95 (2021) 207.
- [16] X.J. Liu, R.L. Lin, B. Jin, Y.B. Zeng, A generalized dressing approach for solving the extended KP and the extended mKP hierarchy, *J. Math. Phys.* 50 (2009) 053506.
- [17] W. Oevel, C. Rogers, Gauge transformations and reciprocal links in  $2 + 1$  dimensions, *Rev. Math. Phys.* 05 (1993) 299–330.
- [18] B.A. Kupershmidt, The quasiclassical limit of the modified KP hierarchy, *J. Phys. A, Math. Gen.* 23 (1990) 871–886.
- [19] J.H. Chang, M.H. Tu, On the Miura Map between the dispersionless KP and dispersionless modified KP hierarchies, *J. Math. Phys.* 41 (2000) 5391.
- [20] Y.T. Chen, M.H. Tu, A note on the Dispersionless Dym Hierarchy, *Lett. Math. Phys.* 63 (2003) 125–139.
- [21] H.X. Wu, Y.B. Zeng, A new extended dispersionless mKP hierarchy and hodograph solution, *Commun. Nonlinear Sci. Numer. Simul.* 17 (2012) 2766–2775.
- [22] H.X. Wu, Y.B. Zeng, J.X. Liu, New extension of dispersionless Harry Dym hierarchy, *J. Nonlinear Math. Phys.* 23 (2016) 383–398.
- [23] T. Xiao, Y.B. Zeng, Bäcklund transformations for the constrained dispersionless hierarchies and dispersionless hierarchies with self-consistent sources, *Inverse Probl.* 22 (2006) 869–879.
- [24] T. Xiao, Y.B. Zeng, The constrained dispersionless mKP hierarchy and the dispersionless mKP hierarchy with self-consistent sources, *Phys. Lett. A* 349 (2006) 128–134.
- [25] H. Aratyn, E. Nissimov, S. Pacheva, Virasoro symmetry of constrained KP hierarchies, *Phys. Lett. A* 228 (1997) 164–175.
- [26] J.S. He, Kelei Tian, A. Foerster, W.X. Ma, Additional symmetries and string equation of the CKP hierarchy, *Lett. Math. Phys.* 81 (2007) 119–134.
- [27] C.Z. Li, J.S. He, Virasoro symmetry of the constrained multicomponent Kadomtsev-Petviashvili hierarchy and its integrable discretization, *Theor. Math. Phys.* 187 (2016) 871–887.
- [28] K.L. Tian, H.J. Song, J.P. Cheng, Y. Cheng, Additional symmetries of constrained CKP and BKP hierarchies, *Sci. China Math.* 54 (2011) 257–268.
- [29] H. Aratyn, Constrained KP hierarchies: additional symmetries, Darboux-Bäcklund solutions and relations to multi-matrix models, *Int. J. Mod. Phys. A* 12 (1999) 1265–1340.
- [30] K.L. Tian, S. Li, G. Yi, Y. Xu, Additional symmetries of the dispersionless cKP hierarchy, *Phys. Lett. B* 822 (2021) 136643.
- [31] T. Takebe, A note on the modified KP hierarchy and its (yet another) dispersionless limit, *Lett. Math. Phys.* 59 (2002) 157–172.
- [32] M.Y. Hu, C.Z. Li, Additional symmetries of the dispersionless extended noncommutative Gelfand-Dickey hierarchy, *Int. J. Geom. Methods Mod. Phys.* 18 (2021) 2150145.